
Bioconversion of Methanol into Value-added Chemicals in Native and Synthetic Methylo-trophs

Min Zhang¹, Xiao-jie Yuan¹, Cong Zhang¹, Li-ping Zhu¹, Xu-hua Mo¹, Wen-jing Chen¹ and Song Yang^{1,2*}

¹School of Life Science, Qingdao Agricultural University, Shandong Province Key Laboratory of Applied Mycology, and Qingdao International Center on Microbes Utilizing Biogas, Qingdao, China.

²Key Laboratory of Systems Bioengineering, Ministry of Education, Tianjin University, Tianjin, China.

*Correspondence: yangsong1209@163.com

<https://doi.org/10.21775/cimb.033.225>

Abstract

Methanol, commercially generated from methane, is a renewable chemical feedstock that is highly soluble, relatively inexpensive, and easy to handle. The concept of native methylo-trophic bacteria serving as whole cell catalysts for production of chemicals and materials using methanol as a feedstock is highly attractive. In recent years, the available omics data for methylo-trophic bacteria, especially for *Methylobacterium extorquens*, the best-characterized model methylo-troph, have provided a solid platform for rational engineering of methylo-trophic bacteria for industrial production. In addition, there is a strong interest in converting the more traditional hetero-trophic production platforms towards the use of single carbon substrates, including methanol, through metabolic engineering. In this chapter, we review the recent progress towards achieving the desired growth and production yields from methanol, by genetically engineered native methylo-trophic strains and by the engineered synthetic methylo-trophs.

production of value-added chemicals. In particular, methanol is an important one-carbon (C1) feedstock that can be generated from either synthesis gas (a mixture of CO and H₂) or from biogas, assuming that large quantities of this feedstock could be produced at relatively low market price (Clomburg *et al.*, 2017; Yang *et al.*, 2018). Methylo-trophic bacteria are a diverse group of microbes that can use reduced C1 compounds such as methanol and methane as sole sources of both energy and carbon (Chistoserdova *et al.*, 2009; Chistoserdova, 2018). Some methylo-trophic bacteria, such as *Methylobacterium extorquens*, *Methylococcus capsulatus*, and *Methylomicrobium buryatense*, show robust growth under laboratory conditions (Bird *et al.*, 1971; Guo and Lidstrom, 2006; de la Torre *et al.*, 2015). The availability of whole genome sequences, combined with biochemical studies (Peyraud *et al.*, 2011; Kalyuzhnaya *et al.*, 2013), have provided important insights into their metabolism that are crucial for enabling engineering these types of microorganism for bulk chemical production. Meantime, as an alternative approach, more traditional model microorganisms such as *Escherichia coli*, *Corynebacterium glutamicum* and *Saccharomyces cerevisiae*, have been genetically modified towards acquiring the ability to utilize methanol for growth,

Introduction

One goal of metabolic engineering is to develop renewable and sustainable alternatives in

with a potential for converting methanol into value-added compounds (Whitaker *et al.*, 2015; Bennett *et al.*, 2018b). In this chapter, we review the current status of the fields of engineering native as well as synthetic methylotrophs and provide examples of successfully engineered strains with a potential in producing commercial value-added chemicals.

Metabolic engineering of native methylotrophs for value-added chemicals production

M. extorquens AM1, a pink-pigmented facultative methylotrophic α -proteobacterium capable of growth on both C1 and multi-carbon compounds, has served for decades as the model organism for studying C1 metabolism (Chistoserdova *et al.*, 2003), including its biotechnological applications (Ochsner *et al.*, 2015). Since the 1960s, it has been reported that *M. extorquens* was successfully employed as a producer of certain products such as amino acids (Sirirote *et al.*, 1988). Based on the current understanding, *M. extorquens* employs three interlocked metabolic cycles for carbon assimilation: the serine cycle, the ethylmalonyl-CoA (EMC) pathway, and the poly-3-hydroxybutyrate (PHB) cycle (Fig. 13.1) (Anthony, 2011; Cui *et al.*, 2016). The main function of the EMC pathway is to regenerate glyoxylate from acetyl-CoA for reincorporating it into the Serine cycle during C1

assimilation (Erb *et al.*, 2007; Peyraud *et al.*, 2009). It has been shown that significant metabolic flux goes through the serine cycle and the EMC pathway when cells are grown on methanol, generating a stable supply of acetyl-CoA, which could serve as a precursor for the production of value-added chemicals in engineered *M. extorquens* AM1 strains (Fig. 13.1) (Peyraud *et al.*, 2011; Fu *et al.*, 2016).

Production of mevalonate and terpenoids

Mevalonate and its derivatives terpenoids are promising bulk chemicals. Terpenoids are used as flavours, fragrances, pharmaceuticals, and biofuels, representing the largest and the most structurally diverse family of chemicals in nature (Zhang *et al.*, 2017). To synthesize considerable amounts of terpenoids, a commonly used strategy in bacteria is to introduce an exogenous mevalonate (MEV) pathway instead of utilizing the native methylerythritol phosphate (MEP) pathway for circumventing the regulatory control of the MEP pathway (Zhao *et al.*, 2013). Thus, pathway engineering for mevalonate production is a feasible first step as a platform for diverse terpenoid synthesis. Although *E. coli* is the primary industrial microbe for the production of mevalonate (Nagai *et al.*, 2018), other engineered hosts have the potential to generate it from alternative carbon feedstocks via synthetic metabolic pathways. Zhu and colleagues




Figure 13.1 Schematic of methanol metabolism in *M. extorquens* and target precursors for synthesis of value added compounds. Pink, grey and blue backgrounds represent the primary oxidation module, the dissimilation module, and the assimilation modules, respectively.

engineered a mevalonate pathway by using acetyl-CoA as the initial precursor in *M. extorquens* AM1, to produce mevalonate from methanol (Zhu *et al.*, 2016). A natural operon (MVH) harbouring the *mvaS* and *mvaE* genes from *Enterococcus faecalis*, as well as an artificial operon (MVH) harbouring the *hmgcs1* gene from *Blattella germanica* and the *tchmgr* gene from *Trypanosoma cruzi* were constructed in a plasmid pCM110. Expression of these two operons resulted in the titres of mevalonate of 56 mg/l and 66 mg/l, respectively, in flask cultivations. Further introduction of the *phaA* gene from *Ralstonia eutropha* into the MVH operon, with the goal of improving the conversion from acetyl-CoA into acetoacetyl-CoA increased the mevalonate titre to 180 mg/l, 3.2-fold higher than that of the natural MVE operon. Further modification of the expression level of the *phaA* gene by regulating the strength of the ribosomal binding site (RBS) resulted in an additional increase in mevalonate production to 215 mg/l (Zhu *et al.*, 2016). Moreover, it has been demonstrated that, under these conditions, the supply of acetyl-CoA was limited in engineered *M. extorquens* AM1. To overcome this limitation, a strategy termed mevalonate sensor-assisted transcriptional regulator engineering (SATRE) was employed, to control metabolic flux redistribution toward increasing acetyl-CoA flux from methanol for mevalonate production (Fig. 13.2) (Liang *et al.*, 2017a). A 60% increase

in mevalonate concentration was further achieved following sensor-based high-throughput screening of a QscR transcriptional regulator library. Finally, a fed-batch fermentation in a 5 l bioreactor yielded a mevalonate concentration of 2.67 g/l, which was equivalent to an overall yield of 0.055 mol acetyl-CoA/mol methanol. In addition, Cui and colleagues utilized atmospheric and room temperature plasma (ARTP) mutagenesis, in combination with adaptive laboratory evolution (ALE), generating a mutant with high methanol tolerance whose cell density was 7.1-fold higher than that of the parent strain in 1.25 M (5%, v/v) methanol (Cui *et al.*, 2018). Accordingly, the mevalonate productivity of the mutant strain carrying a mevalonate synthesis pathway was 65% higher than that of the wild-type strain in methanol fed-batch fermentation.

In a separate study, Sonntag and colleagues introduced a mevalonate pathway from *Myxococcus xanthus*, in combination with RBS optimization of α -humulene and farnesyl pyrophosphate synthases, resulting in the increase of titres of α -humulene from 18 mg/l to 54 mg/l in *M. extorquens* AM1 (Sonntag *et al.*, 2015a). Furthermore, the recombinant *M. extorquens* AM1 was grown in methanol-limited fed-batch conditions, with the addition of dodecane to avoid the accumulation of α -humulene in the medium. An average OD₆₀₀ value of 80–90 were achieved, and final product concentrations were up to 1.65 g/l. This is the highest titre of *de*




Figure 13.2 Biosensor-assisted transcriptional regulator engineering approach to increase mevalonate production in *M. extorquens* AM1. QscR is the regulator of the serine cycle for methanol assimilation (Liang *et al.*, 2017).

novo synthesized α -humulene reported to date, which corresponds to 12% of the maximum theoretical yield. The product titres of mevalonate and α -humulene described in these studies are already in a competitive range compared to the industrial platform strains (Harada *et al.*, 2009), demonstrating the potential of *M. extorquens* as a future platform for the production of high-value terpenoids.

Production of carboxylic acids

3-Hydroxypropionic acid (3-HP) is an important platform chemical, serving as a precursor for a wide range of industrial applications, such as the production of acrylic acid and 1,3-propanediol. In particular, acrylic acid has an annual market exceeding \$10 billion. 3-HP can be further polymerized into poly-3-hydroxypropionate, which is a biodegradable and biocompatible thermoplastic polymer. Recently, Yang and colleagues constructed a malonyl-CoA pathway in *M. extorquens* AM1 that reduced malonyl-CoA to synthesize 3-HP through a bifunctional reductase encoded by the *mcr* gene from *Chloroflexus aurantiacus* (Yang *et al.*, 2017). The engineered strain demonstrated 3-HP production from methanol with initial titre of 6.8 mg/l in shake flask cultivation, which was then improved to 91.0 mg/l by increasing the strength of the promoter and the *mcr* gene copy number, and by decreasing the concentration of sodium ions in the culture medium. Notably, when the cells reached the stationary growth phase, 3-HP content was rapidly reduced to lower than 10.0 mg/l within 10 h. Furthermore, based on the combined results from *in vitro* enzyme assays, ¹³C-labeling analysis, metabolomics and gene knockouts, major routes were identified responsible for the degradation of the 3-HP during this phase of growth. This work provides a solid base for future engineering efforts towards bioconversion of methanol into the economically important product 3-HP.

Recently, Rohde and colleagues introduced (R)-3-hydroxybutyryl-CoA-specific, coenzyme B12-dependent mutases from *Bacillus massiliosenegalensis* JC6 into *M. extorquens* AM1, to convert the intermediate of 3-hydroxybutyryl-CoA towards production of another carboxylic acid, 2-hydroxyisobutyric acid, a building block with a high potential for poly-methylmethacrylate production (Rohde *et al.*, 2017). 2-hydroxyisobutyric acid titres of 2.1 g/l were further achieved in fed-batch

bioreactor without affecting the biomass yield, with the yield of 0.11 g/g from methanol. In addition, Sonntag and colleagues overexpressed *yciA* encoding thioesterases, from *E. coli* in *M. extorquens* AM1, an engineering step designed to generate two dicarboxylic acids from their CoA derivatives involved in the EMC pathway, mesaconic acid (with 70 mg/l yield), and methylsuccinic acid (with 60 mg/l yield), respectively (Sonntag *et al.*, 2014). Further increase in accumulation of mesaconyl-CoA (16-fold) and methylsuccinyl-CoA (6-fold) was achieved by adjusting the concentration of cobalt, a growth limiting factor affecting the activities of ethylmalonyl-CoA and methylmalonyl-CoA mutases, in the culture medium, with a combined titre of the two dicarboxylic acids of 0.65 g/l (Sonntag *et al.*, 2015b). This titre represented a yield of 0.17 g/g methanol. Not only these important achievements shed light on the potential in engineering *M. extorquens* AM1 as an alternative microbial factory for producing common carboxylic acids and uncommon dicarboxylic acids from methanol, but they also illustrate the complex metabolic regulation of methylotrophy in this organism.

Production of 1-butanol

1-Butanol has been proposed to be a better alternative to ethanol as a replacement for gasoline because of its many advantages. For instance, 1-butanol is less soluble in water than ethanol, reducing the possibility of contamination in groundwater, and this trait also makes it feasible to use liquid-liquid extraction for 1-butanol recovery in industrial fermentation. Additionally, 1-butanol is compatible with the current petroleum infrastructure and can be blended with gasoline at high ratios. Although clostridia are the primary industrial microbes for the fermentative production of 1-butanol, clostridia-based industrial fermentation has some limitations, such as formation of by-products, spore formation, and low cell density (Lee *et al.*, 2008). Recently, Hu and colleagues engineered a modified CoA-dependent pathway for 1-butanol production from the precursor crotonyl-CoA, by introducing the *ter* gene from *Treponema denticola* and the *adhE2* from *Clostridium acetobutylicum* into *M. extorquens* AM1 (Hu and Lidstrom, 2014). Although tests for 1-butanol production by cells grown on methanol were negative, the engineered strain was able to generate 1-butanol at 15.2 mg/l when grown on

ethylamine while overexpressing the *croR* gene as a means of increasing the supply of crotonyl-CoA (Hu and Lidstrom, 2014). Adaptive laboratory evolution (ALE) was further employed, resulting in isolation of a mutant with 1-butanol tolerance of up to 0.5%, after 200 generations. The adapted strain was able to produce 25.5 mg/l 1-butanol, representing an 87% increase in production compared with the original strain (Hu *et al.*, 2016). Whole genome sequencing revealed a SNP mutation in the adapted strain, in *kefB*, a gene encoding a potassium/proton antiporter. Analysis of a knockout mutant in this gene confirmed that this mutation was indeed responsible for the increased 1-butanol tolerance. Global metabolomics analysis further uncovered that the pools of multiple key metabolites, including fatty acids, amino acids, disaccharides and carotenoids were either increased or decreased in the adapted strain in response to 1-butanol stress. Overall, this study uncovered an array of cellular responses by *M. extorquens* AM1 to solvent stresses, thus providing a genetic blueprint for further rational design of derivatives of this strain for increased 1-butanol production.

Novel insights into methylotrophy in *Methylobacterium*

Although natural methylotrophs represent promising candidates as new microbial cell factories, further efforts are required to better understand the regulation of methylotrophs in these species and to further develop genetic tools essential for improving the product yields. Recently, Ochsner and colleagues carried out experiments involving hyper-saturating transposon mutagenesis, followed up by high-throughput sequencing of mutant populations (Ochsner *et al.*, 2017). This study uncovered 95 genes that have not been previously known to play a role in methylotrophy (Bringel and Vuilleumier, 2017; Ochsner *et al.*, 2017). Notably, this study uncovered that ribulose-1, 5-bisphosphate (RuBP) functions as a master regulator metabolite in methylotrophic metabolism, by uncovering that *prk*, a gene encoding phosphoribulokinase (Prk), responsible for the production of RuBP was essential for growth on methanol. Prk was ultimately demonstrated to be a co-inducer of QscR, a major regulator of the serine cycle.

In another study, Schada von Borzyskowski and colleagues explored the possibility of converting

the heterotrophic *M. extorquens* AM1 into an autotrophic derivative (Schada von Borzyskowski *et al.*, 2018). They constructed a chassis that utilized methanol solely for energy acquisition while forming the biomass entirely from CO₂. First, they engineered a *M. extorquens* AM1 derivative with deletion in *ffl*, a gene encoding formate tetrahydrofolate ligase, causing the inability to grow on methanol. They then heterologously expressed Rubisco from *Rhodospirillum rubrum*, in order to enable CO₂ fixation. As a result, *M. extorquens* AM1Δ*ffl* expressing the newly built Calvin-Benson-Bassham (CBB) cycle could grow in methanol minimal medium with 5% CO₂, although this strain did not sustain long-lasting growth under this condition. This work is the first example of decoupling energy metabolism and carbon assimilation in *M. extorquens* AM1 and establishing synthetic autotrophy with a heterologously derived CBB cycle. This work also demonstrated the importance of the continuous operation of carbon assimilatory cycles in balancing carbon fluxes between biomass building and draining metabolites from central metabolism into the designed biosynthetic pathways. Taken together, these new contributions offer opportunities for elucidating genotype to phenotype relationships in C₁ assimilation and providing novel insights into its regulation, which will be key in engineering native methylotrophs for biotechnological application in the future.

Constructing synthetic methylotrophs for direct methanol conversion

In recent years, there has been increasing interest in engineering non-methylotrophic bacteria toward the use of C1 compounds, through metabolic engineering (Bennett *et al.*, 2018b). This strategy not only increases our fundamental understanding of methylotrophy, but also provides an opportunity to use established industrial bacteria as platforms for converting methanol into value-added products. The efforts to engineer methanol utilization in traditional platform organisms, both prokaryotic (*E. coli*, *C. glutamicum*; Müller *et al.*, 2015b; Witthoff *et al.*, 2015) and eukaryotic *S. cerevisiae* (Dai *et al.*, 2017) have been recently reported.

Overview of assimilatory C₁ metabolism

Generally, there are two types of methanol dehydrogenase. One is PQQ-dependent Mdh in Gram-negative methylotrophs such as *M. extorquens*, another is NAD-dependent Mdh in Gram-positive *Bacillus methanolicus* (Müller *et al.*, 2015a). At least 14 gene products are necessary for PQQ-dependent Mdh function (Lidstrom, 2006), but NAD-dependent Mdh requires one gene for converting methanol to formaldehyde, and its activity can be greatly enhanced by an activator protein ACT (Anne *et al.*, 2013). Moreover, the PQQ cofactor that is required for PQQ-Mdh is not synthesized by the industrial strains such as *E. coli* or *C. glutamicum*. Therefore, NAD-dependent Mdh has become an obvious choice for implementation of methanol to formaldehyde conversion in *E. coli* and *C. glutamicum* (Fig. 13.3).

For assimilation of formaldehyde into central metabolism, two major pathways in natural methylotrophs have been reported, the ribulose monophosphate (RuMP) and the serine cycles. The RuMP cycle is present in several methylotrophs such as *Methylococcus capsulatus* Bath and *Methylobacillus flagellatus* KT (Strom *et al.*, 1974; Chistoserdova *et al.*, 2000). Formaldehyde is condensed with RuMP by hexulose-6-phosphate synthase (HPS), to generate hexulose-6-phosphate. Hexulose-6-phosphate is further isomerized into fructose 6-phosphate by phosphohexuloisomerase (PHI). Fructose 6-phosphate can be cleaved into C₃ intermediates either via the glycolysis pathway or via the Entner–Doudoroff pathway reactions. The RuMP pathway is quite an efficient route for formaldehyde assimilation in terms of ATP consumption and biomass yield. Per one molecule of pyruvate generated, 1 molecule of ATP and 3 molecules of NAD(P)H are produced.

The serine cycle operates in methylotrophic bacteria such as *M. extorquens* AM1 (Chistoserdova and Lidstrom, 1994). In this cycle, formaldehyde is first oxidized to formate, which is subsequently converted into methylene tetrahydrofolate (THF) via the THF-linked C₁ transfer pathway. Methylene-THF is then condensed with glycine to produce serine, which is subsequently transformed to form C₃ and C₄ intermediates for biomass synthesis. In contrast to the RuMP cycle, the serine cycle leads to the consumption of four molecules of

NAD(P)H and three molecules of ATP to produce one molecule of pyruvate.

Engineering bacteria for methanol conversion

In silico modelling predicted that expression of a NAD-linked Mdh and the two first enzymes of the RuMP cycle, Hps and Phi, into a non-methylotroph such as *E. coli* should be sufficient to allow biomass formation from methanol (Müller *et al.*, 2015b). In order to test this hypothesis, Müller and colleagues heterologously expressed respective genes in *E. coli* (Fig. 13.3) (Müller *et al.*, 2015b). The catalytic activities of several candidate enzymes from different microorganisms such as *B. methanolicus* MGA3, *Lysinibacillus fusiformis*, *M. flagellatus* KT, *M. capsulatus* Bath were compared in cell lysates, and subsequently *in vivo* by measuring formaldehyde formation and assimilation. The three genes originating from *B. methanolicus* MGA3 were found to be the most effective in, respectively, methanol oxidation and formaldehyde conversion by *E. coli*. To further demonstrate how the metabolism of *E. coli* was affected by the introduction of the C1 modules, dynamic labelling experiments using ¹³C-labelled methanol were performed in an *E. coli* strain lacking the *frmA* gene that is responsible for formaldehyde detoxification in native *E. coli*. The incorporation of ¹³C into hexose 6-phosphates was measured to be within approximately 40%. Other metabolites of the RuMP cycle, such as ribose 5-phosphate and ribulose 5-phosphate, as well as intermediates of the glycolysis pathway, such as dihydroxyacetone phosphate and phosphoenolpyruvate were detected as well, demonstrating that at least several turnovers have occurred through the synthetic RuMP cycle. However, no methanol-dependent growth of *E. coli* was achieved. Meantime, a similar engineering scheme has also been applied to *C. glutamicum*, demonstrating utilization of methanol as a co-carbon source in a glucose-based medium in (Lessmeier *et al.*, 2015; Witthoff *et al.*, 2015).

Since Mdh from *B. methanolicus* has low affinity toward methanol (K_m of 200 mM), Whitaker and colleagues employed an alternative enzyme from *B. stearothermophilus*, possessing a K_m of 20 mM and V_{max} of 2.1 U/mg for methanol (Price *et al.*, 2016; Whitaker *et al.*, 2017). They engineered an *E. coli* strain expressing *B. stearothermophilus* Mdh




Figure 13.3 Methanol metabolism in synthetic methylotrophs. Blue solid lines represent reactions mediated by the heterologously expression genes. Red solid lines represent gene knockouts. Enzymes (genes): methanol dehydrogenase (*mdh*), hexulose phosphate synthase (*hps*), phosphohexuloisomerase (*phi*), phosphoglucosomerase (*pgi*), formaldehyde dehydrogenase (*frmA*), phosphofructokinase (*pfk*), fructose-bisphosphate aldolase (*fba*), transketolase (*tkt*), ribulose phosphate epimerase (*rpe*), sedoheptulose bisphosphatase (*glpX*), fructose bisphosphatase (*fbp*), ribose phosphate isomerase (*rpi*), glucose-6-phosphate dehydrogenase (*zwf*). Metabolites: glucose 6-phosphate (G6P), 6-phosphogluconolactone (GL6P), 6-phosphogluconate (6PG), ribulose 5-phosphate (Ru5P), ribose 5-phosphate (R5P), hexulose 6-phosphate (H6P), fructose 6-phosphate (F6P), fructose-1, 6-bisphosphate (FBP), dihydroxyacetone phosphate (DHAP), glyceraldehyde 3-phosphate (GAP), xylulose 5-phosphate (X5P), erythrose 4-phosphate (E4P), sedoheptulose bisphosphate (SBP), sedoheptulose 7-phosphate (S7P), 1,3-bisphosphoglycerate (1,3-BPG), 3-phosphoglycerate (3-PG), 2-phosphoglycerate (2-PG), phosphoenolpyruvate (PEP)

and *B. methanolicus* Hps and Phi, and found that small amounts of supplements such as yeast extract enabled initiation of growth of *E. coli* on methanol. When the engineered *E. coli* strain was grown in M9 minimal medium with 1 g/l yeast extract and 60 mM methanol, a cell biomass increase by 30% was observed, corresponding to biomass yield of 0.289 gCDW/g methanol. This yield is close to biomass yields reported for native methylotrophs. ^{13}C labelling analysis demonstrated that not only the RuMP cycle intermediates were labelled, but

also the metabolites in the TCA cycle and related amino acids, supporting biomass generation from methanol. Moreover, in this study, a 65% improvement in flavonoid production was observed in the presence of 60 mM methanol. Additional engineering effort included heterologous expression of the non-oxidative pentose phosphate pathway (PPP) consisting of five enzymes (ribulose phosphate epimerase (*rpe*), fructose-bisphosphate aldolase (*fba*), sedoheptulose bisphosphatase (*glpX*), phosphofructokinase (*pfk*) and transketolase (*tkt*)

(Bennet *et al.*, 2018a). The introduction of the five genes, all originating from *B. methanoli*, resulted in increased flux through the PPP pathway, further enhancing methanol assimilation by the engineered *E. coli* (Fig. 13.3) (Bennett *et al.*, 2018a). In addition, Bennett and colleagues deleted the phosphoglucosomerase gene (*pgi*), further increasing carbon flux through the oxidative PPP, for increased supply of Ru5P. This change led to a significant improvement of methanol assimilation and to increased acetone titres during co-culturing on methanol and glucose.

In order to further increase methanol consumption, Meyer and colleagues used genetic constraints to develop an approach to link methanol assimilation to growth supported by a multi-carbon source (Meyer *et al.*, 2018). Via *in silico* analysis of a metabolic scenario including the presence of the *mdh-hps-phi* operon, they predicted that the deletion of *edd*, encoding phosphoglucuronate dehydratase, and *rpiAB*, encoding ribose-5-phosphate isomerase, would result in failure to grow on gluconate, but this variant would grow in the presence of methanol. ALE was then employed to improve co-consumption of methanol and gluconate. After 35 generations, the adapted strain showed 3-fold increase in final OD₆₀₀, compared with the initially engineered strain. Furthermore, deletion of the NAD-dependent malate dehydrogenase (*maldh*), which is involved in the TCA cycle, was predicted to increase the ratio of NAD⁺/NADH, potentially increasing methanol oxidation and methanol dependency. After additional rounds of ALE of the $\Delta edd \Delta rpiAB \Delta maldh$ variant of *E. coli*, growth rate was further improved, from 0.017/h to 0.081/h when during co-cultivated on methanol and gluconate. This evolved strain had a methanol consumption rate of 13 mmol/gCDW/h, which is comparable with the reported rate for *M. extorquens* AM1. This work provided an important basis for further optimizing synthetic methylotrophic organisms, in the future.

Engineering a formaldehyde-inducible promoter for improved growth on methanol

In synthetic methylotrophs, methanol is converted to the toxic intermediate formaldehyde by Mdh, catalyzing the initial reaction of methanol conversion. *E. coli* has a native formaldehyde-inducible promoter, *Pfrm*, which negatively regulates the

formaldehyde detoxification operon *frmRAB* (Fig. 13.4) (Danilov and Ismailov, 1989). Recently, Rohlhill and Sandoval constructed a library of mutant variants of *Pfrm* and used the combination of fluorescence-activated cell sorting and high-throughput sequencing (sort-seq) to screen formaldehyde-responsive promoters (Rohlhill and Sandoval, 2017). Two 19 base pair repressor binding sites, one upstream of the -35 RNA polymerase (RNAP) binding site and one overlapping with the -10 site were identified and subsequently engineered, resulting in 13-fold higher expression compared with the native promoter. They also found that the mutant *Pfrm*, placed upstream of the methanol assimilation module (i.e. the *mdh-hps-phi* operon), resulted in increased biomass from methanol, likely due to the improved balance in formaldehyde production/consumption. Further, Woolston and colleagues rationally designed the sequence between -10 and -35 region of the *Pfrm*, and found that the engineered promoter possessed increased sensitivity and wider dynamic range of sensing formaldehyde, at levels as low as 1 μ M. This synthetic biosensor was also employed for optimizing the expression levels of *mdh* and *hps/phi*, to maximize the methanol assimilation flux (Woolston *et al.*, 2018).

Preliminary test for direct methanol utilization in yeast

Besides prokaryotic methylotrophs, *S. cerevisiae* also presents a great potential as a microbial chassis for methanol conversion, due to its robust tolerance to high concentrations of methanol. Dai and colleagues integrated a methanol oxidation pathway including alcohol oxidases (AOX), catalases (CAT), dihydroxyacetone synthases (DAS1/2), and dihydroxyacetone kinase (DAK), originating from the methylotrophic yeast *Pichia pastoris*, into the chromosome of *S. cerevisiae* (Dai *et al.*, 2017). These methylotrophic enzymes with targeting signals from *P. pastoris* were found to be functionally expressed and located in the peroxisomes in the recombinant *S. cerevisiae*. The engineered *S. cerevisiae* showed 1.04 g/l consumption of methanol and 3.13% increase in cell growth, accompanied with 0.26 g/l of pyruvate production in the fermentation broth when using methanol as the sole carbon source. In another study, *P. pastoris* was engineered to produce lovastatin and its precursor monacolin




Figure 13.4 Regulation of the *frm* operon. (A) FrmR binds to the promoter (*Pfrm*) upstream of the *frmRAB* gene cluster, repressing transcription. (B) FrmR is conformationally modified by binding formaldehyde, permitting transcription.

J from methanol (Liu *et al.*, 2018a,b). A co-culture of *P. pastoris* was optimized for bioreactor fermentations with the yields of 593.9 mg/l monacolin J and 250.8 mg/l lovastatin. Although much further work is required to improve methanol utilization and concomitant production of value-added chemicals, these preliminary studies demonstrate a potential in engineering non-methylotrophic eukaryotic microbes for methanol utilization and scaling their use toward industrial biotechnological processes in the future.

Conclusions

In the recent years, native methylotrophs such as *M. extorquens* AM1 have gained interest highlighting the important role of methanol as a future alternative carbon feedstock. With understanding of native metabolic pathways and further development of metabolic engineering tools in *M. extorquens* AM1, it may be feasible to rationally manipulate this microorganism for producing bio-based chemicals and biofuels (Lieven *et al.*, 2018). However, further strain optimization is required to make this system industrially relevant. Alternatively, engineering synthetic methylotrophic bacteria and yeasts provide attractive alternative platforms for methanol conversion. Examples of methanol assimilation by

engineered synthetic methylotrophs demonstrate the potential for using traditional industrial micro-organism as chassis for production of value-added chemicals from methanol.

Future trends

Two key roadblocks were identified while attempting to engineer native methylotrophs for methanol conversion. First is the insufficient knowledge on the regulatory mechanisms. For example, engineering even one of the best-studied methylotrophs, *M. extorquens*, for biotechnological applications requires more knowledge on both the essential genes and their associated regulatory networks. The integrated multiple 'omics' approaches and the genome editing tools that are becoming available for a multitude of organisms, will aid in elucidating the essential sets of proteins-metabolites that control the regulatory networks (Jiang *et al.*, 2013; Diether and Sauer, 2017). Recent discovery of the phosphoribulokinase gene playing an unexpected but essential regulatory role provides one example (Ochsner *et al.*, 2017). The second roadblock is the lack of well-developed genetic tools and genetic elements. Schada von Borzyskowski and collaborators demonstrated that extensive sets of plasmids with different properties are required to allow for

efficient construction of synthetic operons with different promoter strengths in *M. extorquens* AM1 (Schada von Borzyskowski *et al.*, 2015). Liang and colleagues developed biosensor-assisted transcriptional regulator engineered to control metabolic flux re-distribution between C1 assimilation and chemicals biosynthesis (Liang *et al.*, 2017a). Future work is necessary to explore regulatory mechanism that may not be directly involved in methanol metabolism, but could potentially improve methanol assimilation. It is also necessary to develop advanced engineering tools, such as promoters and RBS's of different strengths to allow for fine tuning gene expression, as well as the convenient Cre/loxP systems for integrating entire heterologous pathways into the genomes of *M. extorquens* (Liang *et al.*, 2017b) and other potential platforms.

For synthetic methylotrophs, a persisting challenge remains the lack of growth on methanol as the sole carbon and energy source. One limitation is the efficiency of methanol oxidation catalyzed by NAD-dependent Mdh. So far, the most studied is the Mdh from *B. methanolicus*, which exhibits low enzyme specificity toward methanol and dependency on an activator protein (Price *et al.*, 2016). Recently, Wu and collaborators characterized a NAD-dependent alcohol dehydrogenase (Mdh2) from *Cupriavidus necator* that does not require an activator (Wu *et al.*, 2016). Further, they engineered a variant of Mdh2 by using directed molecular evolution that showed a 6-fold increase in K_{cat}/K_m for methanol and 10-fold decrease in K_{cat}/K_m for 1-butanol than that of Mdh from *B. methanolicus*. Another limitation is the insufficient flux toward regeneration of Ru5P. Bennett and colleagues demonstrated significant progress towards solving this problem, but further progress is needed (Bennett *et al.*, 2018a). Recently, Cui and colleagues used ARTP mutagenesis to identify a mutant with obvious improvement of methanol tolerance and consumption (Cui *et al.*, 2018). This state-of-the-art technique could also be adapted to evolve current synthetic methylotrophs and screen mutants for increased methanol conversion in the future.

Acknowledgements

Grant information: This work was supported by the National Natural Science Foundation of China (Grant No. 21776149) and Shandong Provincial Key Research and Development Project, China (2016GSF117026).

References

- Anne, K., Heggeset, T.M.B., Müller, J.E.N., Kupper, C.E., Olha, S., Vorholt, J.A., Ellingsen, T.E., and Trygve, B. (2013). Methylotrophic *Bacillus methanolicus* encodes two chromosomal and one plasmid Born NAD⁺-dependent methanol dehydrogenase paralogs with different catalytic and biochemical properties. *PLOS ONE* 8, e59188.
- Anthony, C. (2011). How half a century of research was required to understand bacterial growth on C1 and C2 compounds; the story of the serine cycle and the ethylmalonyl-CoA pathway. *Sci. Prog.* 94, 109–137.
- Bennett, R.K., Gonzalez, J.E., Whitaker, W.B., Antoniewicz, M.R., and Papoutsakis, E.T. (2018a). Expression of heterologous non-oxidative pentose phosphate pathway from *Bacillus methanolicus* and phosphoglucose isomerase deletion improves methanol assimilation and metabolite production by a synthetic *Escherichia coli* methylotroph. *Metab. Eng.* 45, 75–85.
- Bennett, R.K., Steinberg, L.M., Chen, W., and Papoutsakis, E.T. (2018b). Engineering the bioconversion of methane and methanol to fuels and chemicals in native and synthetic methylotrophs. *Curr. Opin. Biotechnol.* 50, 81–93.
- Bird, C.W., Lynch, J.M., Pirt, F.J., and Reid, W.W. (1971). Steroids and squalene in *Methylococcus capsulatus* grown on methane. *Nature* 230, 473–474.
- Bringel, F., and Vuilleumier, S. (2017). Metabolic regulation: a master role for ribulose-1,5-bisphosphate in one-carbon assimilation. *Curr. Biol.* 27, R1127–R1129.
- Chistoserdova, L. (2018). Applications of methylotrophs: can single carbon be harnessed for biotechnology? *Curr. Opin. Biotechnol.* 50, 189–194.
- Chistoserdova, L.V., and Lidstrom, M.E. (1994). Genetics of the serine cycle in *Methylobacterium extorquens* AM1: identification of *sgaA* and *mtdA* and sequences of *sgaA*, *hprA*, and *mtdA*. *J. Bacteriol.* 176, 1957–1968.
- Chistoserdova, L., Gomelsky, L., Vorholt, J.A., Gomelsky, M., Tsygankov, Y.D., and Lidstrom, M.E. (2000). Analysis of two formaldehyde oxidation pathways in *Methylobacillus flagellatus* KT, a ribulose monophosphate cycle methylotroph. *Microbiology* 146, 233–238. <https://doi.org/10.1099/00221287-146-1-233>
- Chistoserdova, L., Chen, S.W., Lapidus, A., and Lidstrom, M.E. (2003). Methylotrophy in *Methylobacterium extorquens* AM1 from a genomic point of view. *J. Bacteriol.* 185, 2980–2987.
- Chistoserdova, L., Kalyuzhnaya, M.G., and Lidstrom, M.E. (2009). The expanding world of methylotrophic metabolism. *Annu. Rev. Microbiol.* 63, 477–499. <https://doi.org/10.1146/annurev.micro.091208.073600>
- Clomburg, J.M., Crumbley, A.M., and Gonzalez, R. (2017). Industrial biomanufacturing: The future of chemical production. *Science* 355, aag0804.
- Cui, J., Good, N.M., Hu, B., Yang, J., Wang, Q., Sadilek, M., and Yang, S. (2016). Metabolomics revealed an association of metabolite changes and defective growth in *Methylobacterium extorquens* AM1 overexpressing *ecm* during growth on methanol. *PLOS ONE* 11, e0154043. <https://doi.org/10.1371/journal.pone.0154043>
- Cui, L.Y., Wang, S.S., Guan, C.G., Liang, W.F., Xue, Z.L., Zhang, C., and Xing, X.H. (2018). Breeding of

- methanol-tolerant *Methylobacterium extorquens* AM1 by atmospheric and room temperature plasma mutagenesis combined with adaptive laboratory evolution. *Biotechnol. J.* 13, e1700679. <https://doi.org/10.1002/biot.201700679>
- Dai, Z., Gu, H., Zhang, S., Xin, F., Zhang, W., Dong, W., Ma, J., Jia, H., and Jiang, M. (2017). Metabolic construction strategies for direct methanol utilization in *Saccharomyces cerevisiae*. *Bioresour. Technol.* 245, 1407–1412.
- Danilov, V.S., and Ismailov, A.D. (1989). Bacterial luciferase as a biosensor of biologically active compounds. *Biotechnology* 11, 39–78.
- de la Torre, A., Metivier, A., Chu, F., Laurens, L.M., Beck, D.A., Pienkos, P.T., Lidstrom, M.E., and Kalyuzhnaya, M.G. (2015). Genome-scale metabolic reconstructions and theoretical investigation of methane conversion in *Methylobacterium buryatense* strain 5G(B1). *Microb. Cell Fact.* 14, 188. <https://doi.org/10.1186/s12934-015-0377-3>
- Diether, M., and Sauer, U. (2017). Towards detecting regulatory protein-metabolite interactions. *Curr. Opin. Microbiol.* 39, 16–23.
- Erb, T.J., Berg, I.A., Brecht, V., Müller, M., Fuchs, G., and Alber, B.E. (2007). Synthesis of C5-dicarboxylic acids from C2-units involving crotonyl-CoA carboxylase/reductase: the ethylmalonyl-CoA pathway. *Proc. Natl. Acad. Sci. U.S.A.* 104, 10631–10636.
- Fu, Y., Beck, D.A., and Lidstrom, M.E. (2016). Difference in C3-C4 metabolism underlies tradeoff between growth rate and biomass yield in *Methylobacterium extorquens* AM1. *BMC Microbiol.* 16, 156. <https://doi.org/10.1186/s12866-016-0778-4>
- Guo, X., and Lidstrom, M.E. (2006). Physiological analysis of *Methylobacterium extorquens* AM1 grown in continuous and batch cultures. *Arch. Microbiol.* 186, 139–149. <https://doi.org/10.1007/s00203-006-0131-7>
- Harada, H., Yu, F., Okamoto, S., Kuzuyama, T., Utsumi, R., and Misawa, N. (2009). Efficient synthesis of functional isoprenoids from acetoacetate through metabolic pathway-engineered *Escherichia coli*. *Appl. Microbiol. Biotechnol.* 81, 915–925. <https://doi.org/10.1007/s00253-008-1724-7>
- Hu, B., and Lidstrom, M.E. (2014). Metabolic engineering of *Methylobacterium extorquens* AM1 for 1-butanol production. *Biotechnol. Biofuels* 7, 156. <https://doi.org/10.1186/s13068-014-0156-0>
- Hu, B., Yang, Y.M., Beck, D.A., Wang, Q.W., Chen, W.J., Yang, J., Lidstrom, M.E., and Yang, S. (2016). Comprehensive molecular characterization of *Methylobacterium extorquens* AM1 adapted for 1-butanol tolerance. *Biotechnol. Biofuels* 9, 84. <https://doi.org/10.1186/s13068-016-0497-y>
- Jiang, W., Bikard, D., Cox, D., Zhang, F., and Marraffini, L.A. (2013). RNA-guided editing of bacterial genomes using CRISPR-Cas systems. *Nat. Biotechnol.* 31, 233–239. <https://doi.org/10.1038/nbt.2508>
- Kalyuzhnaya, M.G., Yang, S., Rozova, O.N., Smalley, N.E., Clubb, J., Lamb, A., Gowda, G.A., Raftery, D., Fu, Y., Bringel, F., et al. (2013). Highly efficient methane biocatalysis revealed in a methanotrophic bacterium. *Nat. Commun.* 4, 2785. <https://doi.org/10.1038/ncomms3785>
- Lee, S.Y., Park, J.H., Jang, S.H., Nielsen, L.K., Kim, J., and Jung, K.S. (2008). Fermentative butanol production by Clostridia. *Biotechnol. Bioeng.* 101, 209–228. <https://doi.org/10.1002/bit.22003>
- Leßmeier, L., Pfeifenschneider, J., Carnicer, M., Heux, S., Portais, J.C., and Wendisch, V.F. (2015). Production of carbon-13-labeled cadaverine by engineered *Corynebacterium glutamicum* using carbon-13-labeled methanol as co-substrate. *Appl. Microbiol. Biotechnol.* 99, 10163–10176. <https://doi.org/10.1007/s00253-015-6906-5>
- Liang, W.F., Cui, L.Y., Cui, J.Y., Yu, K.W., Yang, S., Wang, T.M., Guan, C.G., Zhang, C., and Xing, X.H. (2017a). Biosensor-assisted transcriptional regulator engineering for *Methylobacterium extorquens* AM1 to improve mevalonate synthesis by increasing the acetyl-CoA supply. *Metab. Eng.* 39, 159–168.
- Liang, W.F., Sun, M.Y., Cui, L.Y., Zhang, C., and Xing, X.H. (2017b). Cre/loxP-mediated multicopy integration of the mevalonate operon into the genome of *Methylobacterium extorquens* AM1. *Appl. Biochem. Biotechnol.* 2, 1–13.
- Lidstrom, M.E. (2006). Aerobic Methylotrophic Prokaryotes. In *The Prokaryotes*, Vol. 2: Ecophysiology and Biochemistry, Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K.-H., and Stackebrandt, E., eds. (Springer-Verlag, New York, NY), pp. 618–634.
- Lieven, C., Herrgård, M.J., and Sonnenschein, N. (2018). Microbial Methylotrophic metabolism: recent metabolic modeling efforts and their applications in industrial biotechnology. *Biotechnol. J.* 13, e1800011. <https://doi.org/10.1002/biot.201800011>
- Liu, Y., Bai, C., Xu, Q., Yu, J., Zhou, X., Zhang, Y., and Cai, M. (2018a). Improved methanol-derived lovastatin production through enhancement of the biosynthetic pathway and intracellular lovastatin efflux in methylotrophic yeast. *Bioresources & Bioprocessing* 5, 22.
- Liu, Y., Tu, X., Xu, Q., Bai, C., Kong, C., Liu, Q., Yu, J., Peng, Q., Zhou, X., Zhang, Y., et al. (2018b). Engineered monoculture and co-culture of methylotrophic yeast for de novo production of monacolin J and lovastatin from methanol. *Metab. Eng.* 45, 189–199.
- Meyer, F., Keller, P., Hartl, J., Gröninger, O.G., Kiefer, P., and Vorholt, J.A. (2018). Methanol-essential growth of *Escherichia coli*. *Nat. Commun.* 9, 1508. <https://doi.org/10.1038/s41467-018-03937-y>
- Müller, J.E., Meyer, F., Litsanov, B., Kiefer, P., and Vorholt, J.A. (2015a). Core pathways operating during methylotrophy of *Bacillus methanolicus* MGA3 and induction of a bacillithiol-dependent detoxification pathway upon formaldehyde stress. *Mol. Microbiol.* 98, 1089–1100. <https://doi.org/10.1111/mmi.13200>
- Müller, J.E.N., Meyer, F., Litsanov, B., Kiefer, P., Potthoff, E., Heux, S., Quax, W.J., Wendisch, V.F., Brautaset, T., Portais, J.C., et al. (2015b). Engineering *Escherichia coli* for methanol conversion. *Metab. Eng.* 28, 190–201.
- Nagai, H., Masuda, A., Toya, Y., Matsuda, F., and Shimizu, H. (2018). Metabolic engineering of mevalonate-producing *Escherichia coli* strains based on thermodynamic analysis. *Metab. Eng.* 47, 1–9.
- Ochsner, A.M., Sonntag, F., Buchhaupt, M., Schrader, J., and Vorholt, J.A. (2015). *Methylobacterium extorquens*:

- methylotrophy and biotechnological applications. *Appl. Microbiol. Biotechnol.* 99, 517–534. <https://doi.org/10.1007/s00253-014-6240-3>
- Ochsner, A.M., Christen, M., Hemmerle, L., Peyraud, R., Christen, B., and Vorholt, J.A. (2017). Transposon sequencing uncovers an essential regulatory function of phosphoribulokinase for methylotrophy. *Curr. Biol.* 27, 2579–2588.e6.
- Peyraud, R., Kiefer, P., Christen, P., Massou, S., Portais, J.C., and Vorholt, J.A. (2009). Demonstration of the ethylmalonyl-CoA pathway by using ¹³C metabolomics. *Proc. Natl. Acad. Sci. U.S.A.* 106, 4846–4851. <https://doi.org/10.1073/pnas.0810932106>
- Peyraud, R., Schneider, K., Kiefer, P., Massou, S., Vorholt, J.A., and Portais, J.C. (2011). Genome-scale reconstruction and system level investigation of the metabolic network of *Methylobacterium extorquens* AM1. *BMC Syst. Biol.* 5, 189. <https://doi.org/10.1186/1752-0509-5-189>
- Price, J.V., Chen, L., Whitaker, W.B., Papoutsakis, E., and Chen, W. (2016). Scaffoldless engineered enzyme assembly for enhanced methanol utilization. *Proc. Natl. Acad. Sci. U.S.A.* 113, 12691–12696. <https://doi.org/10.1073/pnas.1601797113>
- Rohde, M.T., Tischer, S., Harms, H., and Rohwerder, T. (2017). Production of 2-hydroxyisobutyric acid from methanol by *Methylobacterium extorquens* AM1 Expressing (R)-3-Hydroxybutyryl Coenzyme A-Isomerizing Enzymes. *Appl. Environ. Microbiol.* 83, e02622–16.
- Rohllhill, J., Sandoval, N.R., and Papoutsakis, E.T. (2017). Sort-seq approach to engineering a formaldehyde-inducible promoter for dynamically regulated *Escherichia coli* growth on methanol. *ACS Synth. Biol.* 6, 1584–1595. <https://doi.org/10.1021/acssynbio.7b00114>
- Schada von Borzyskowski, L., Remus-Emsermann, M., Weishaupt, R., Vorholt, J.A., and Erb, T.J. (2015). A set of versatile brick vectors and promoters for the assembly, expression, and integration of synthetic operons in *Methylobacterium extorquens* AM1 and other alphaproteobacteria. *ACS Synth. Biol.* 4, 430–443. <https://doi.org/10.1021/sb500221v>
- Schada von Borzyskowski, L., Carrillo, M., Leupold, S., Glatter, T., Kiefer, P., Weishaupt, R., Heinemann, M., and Erb, T.J. (2018). An engineered Calvin-Benson-Bassham cycle for carbon dioxide fixation in *Methylobacterium extorquens* AM1. *Metab. Eng.* 47, 423–433.
- Sirirote, P., Tsuneo, Y., and Shoichi, S. (1988). L-serine production from methanol and glycine with an immobilized methylotroph. *J. Ferment. Technol.* 66, 291–297.
- Sonntag, F., Buchhaupt, M., and Schrader, J. (2014). Thioesterases for ethylmalonyl-CoA pathway derived dicarboxylic acid production in *Methylobacterium extorquens* AM1. *Appl. Microbiol. Biotechnol.* 98, 4533–4544. <https://doi.org/10.1007/s00253-013-5456-y>
- Sonntag, F., Kroner, C., Lubuta, P., Peyraud, R., Horst, A., Buchhaupt, M., and Schrader, J. (2015a). Engineering *Methylobacterium extorquens* for de novo synthesis of the sesquiterpenoid α -humulene from methanol. *Metab. Eng.* 32, 82–94.
- Sonntag, F., Müller, J.E., Kiefer, P., Vorholt, J.A., Schrader, J., and Buchhaupt, M. (2015b). High-level production of ethylmalonyl-CoA pathway-derived dicarboxylic acids by *Methylobacterium extorquens* under cobalt-deficient conditions and by polyhydroxybutyrate negative strains. *Appl. Microbiol. Biotechnol.* 99, 3407–3419. <https://doi.org/10.1007/s00253-015-6418-3>
- Strom, T., Ferenci, T., and Quayle, J.R. (1974). The carbon assimilation pathways of *Methylococcus capsulatus*, *Pseudomonas methanica* and *Methylosinus trichosporium* (OB3B) during growth on methane. *Biochem. J.* 144, 465–476.
- Whitaker, W.B., Sandoval, N.R., Bennett, R.K., Fast, A.G., and Papoutsakis, E.T. (2015). Synthetic methylotrophy: engineering the production of biofuels and chemicals based on the biology of aerobic methanol utilization. *Curr. Opin. Biotechnol.* 33, 165–175. <https://doi.org/10.1016/j.copbio.2015.01.007>
- Whitaker, W.B., Jones, J.A., Bennett, R.K., Gonzalez, J.E., Vernacchio, V.R., Collins, S.M., Palmer, M.A., Schmidt, S., Antoniewicz, M.R., Koffas, M.A., et al. (2017). Engineering the biological conversion of methanol to specialty chemicals in *Escherichia coli*. *Metab. Eng.* 39, 49–59.
- Witthoff, S., Schmitz, K., Niedenführ, S., Nöh, K., Noack, S., Bott, M., and Marienhagen, J. (2015). Metabolic engineering of *Corynebacterium glutamicum* for methanol metabolism. *Appl. Environ. Microbiol.* 81, 2215–2225. <https://doi.org/10.1128/AEM.03110-14>
- Woolston, B.M., Roth, T., Kohale, I., Liu, D.R., and Stephanopoulos, G. (2018). Development of a formaldehyde biosensor with application to synthetic methylotrophy. *Biotechnol. Bioeng.* 115, 206–215. <https://doi.org/10.1002/bit.26455>
- Wu, T.Y., Chen, C.T., Liu, J.T., Bogorad, I.W., Damoiseaux, R., and Liao, J.C. (2016). Characterization and evolution of an activator-independent methanol dehydrogenase from *Cupriavidus necator* N-1. *Appl. Microbiol. Biotechnol.* 100, 4969–4983. <https://doi.org/10.1007/s00253-016-7320-3>
- Yang, J., Zhang, C.T., Yuan, X.J., Zhang, M., Mo, X.H., Tan, L.L., Zhu, L.P., Chen, W.J., Yao, M.D., Hu, B., Yang, S. (2018). Metabolic engineering of *Methylobacterium extorquens* AM1 for the production of butadiene precursor. *Microb Cell Fact.* 17, 194. doi: 10.1186/s12934-018-1042-4.
- Yang, Y.M., Chen, W.J., Yang, J., Zhou, Y.M., Hu, B., Zhang, M., Zhu, L.P., Wang, G.Y., and Yang, S. (2017). Production of 3-hydroxypropionic acid in engineered *Methylobacterium extorquens* AM1 and its reassimilation through a reductive route. *Microb. Cell Fact.* 16, 179. <https://doi.org/10.1186/s12934-017-0798-2>
- Zhang, L., Xiao, W.H., Wang, Y., Yao, M.D., Jiang, G.Z., Zeng, B.X., Zhang, R.S., and Yuan, Y.J. (2017). Chassis and key enzymes engineering for monoterpenes production. *Biotechnol. Adv.* 35, 1022–1031.
- Zhao, L., Chang, W.C., Xiao, Y., Liu, H.W., and Liu, P. (2013). Methylerythritol phosphate pathway of isoprenoid biosynthesis. *Annu. Rev. Biochem.* 82, 497–530. <https://doi.org/10.1146/annurev-biochem-052010-100934>
- Zhu, W.L., Cui, J.Y., Cui, L.Y., Liang, W.F., Yang, S., Zhang, C., and Xing, X.H. (2016). Bioconversion of methanol to value-added mevalonate by engineered *Methylobacterium extorquens* AM1 containing an optimized mevalonate pathway. *Appl. Microbiol. Biotechnol.* 100, 2171–2182. <https://doi.org/10.1007/s00253-015-7078-z>