

Mobile Genetic Elements in Streptococci

Miao Lu[#], Tao Gong[#], Anqi Zhang, Boyu Tang, Jiamin Chen, Zhong Zhang, Yuqing Li^{*}, Xuedong Zhou^{*}

State Key Laboratory of Oral Diseases, National Clinical Research Center for Oral Diseases, West China Hospital of Stomatology, Sichuan University, Chengdu, PR China.

[#]Miao Lu and Tao Gong contributed equally to this work.

^{*}Address correspondence to: liyuqing@scu.edu.cn, zhouxd@scu.edu.cn

Abstract

Streptococci are a group of Gram-positive bacteria belonging to the family Streptococcaceae, which are responsible of multiple diseases. Some of these species can cause invasive infection that may result in life-threatening illness. Moreover, antibiotic-resistant bacteria are considerably increasing, thus imposing a global consideration. One of the main causes of this resistance is the horizontal gene transfer (HGT), associated to gene transfer agents including transposons, integrons, plasmids and bacteriophages. These agents, which are called mobile genetic elements (MGEs), encode proteins able to mediate DNA movements. This review briefly describes MGEs in streptococci, focusing on their structure and properties related to HGT and antibiotic resistance.

Introduction

Streptococci are a group of Gram-positive bacteria widely distributed across human and animals. Unlike the *Staphylococcus* species, streptococci are catalase negative and are subclassified into the three subspecies alpha, beta and gamma according to the partial, complete or absent hemolysis induced, respectively. The beta hemolytic streptococci species are further classified by the cell wall carbohydrate composition (Lancefield, 1933) and according to human diseases in Lancefield groups A, B, C and G. group C and group G antigen are relatively rare compared with group A and group B antigen (Korona-Glowniak et al., 2015). Streptococci cause numerous diseases in human and animals, and many of them can cause serious and invasive infections damaging public health.

Group A streptococcus (GAS), also known as *Streptococcus pyogenes*, causes serious infections such as streptococcal toxic shock syndrome necrotizing fasciitis, and causes mild human infections such as impetigo and pharyngitis. Repeated GAS infections can cause autoimmune diseases, such as acute rheumatic fever, rheumatic heart disease and acute post-streptococcal glomerulonephritis (Parks et al., 2015). Group B streptococcus (GBS), also known as *Streptococcus agalactiae*, is included in the normal flora of the rectum and female urogenital tract, which under normal circumstances causes no harm, but it may trigger infection when the body defense system is weak. The infection can be transferred to a neonate from the mother colonized with GBS, causing meningitis and sepsis in newborns (Verani et al., 2010). The most represented specie belonging to the groups C and G streptococci is the *Streptococcus equisimilis*, a subspecies of *Streptococcus dysgalactiae*, which is usually present in human upper airways

(Jensen and Kilian, 2012). *S. dysgalactiae* is able to colonize other sites like skin, causing superficial and invasive infections such as septic arthritis, skin and tissue infections (Brandt and Spellerberg, 2009). *Streptococcus pneumoniae* is part of the normal flora of the human upper respiratory tract. The primary site of pneumococcal colonization in human body is the nasopharynx. It can cause a wide spectrum of infections, including invasive and non-invasive diseases. Pneumococcal infection can occur at multiple sites including the meninges and lung parenchyma, and is the primary cause of community acquired pneumonia (CAP) and a main cause of bacteremia, especially in infants and older adults (Remington and Sligl, 2014). The viridans streptococci, belonging to a heterogeneous group, are commensal flora of the oral cavity, upper respiratory, gastrointestinal and female genital tracts. These species can be divided into five groups named *Anginosus*, *Mutans*, *Mitis*, *Salivarius* and *Sanguinis* (Facklam, 2002). The viridians streptococci can cause severe infections including infective endocarditis, dental caries and deep seated abscesses (Menon, 2016). The *Streptococcus anginosus* group, previously known as *Streptococcus milleri*, includes three distinct species: *Streptococcus intermedius*, *Streptococcus anginosus* and *Streptococcus constellatus* (Claridge et al., 2001). These species, constituting the normal flora of the gastrointestinal tract and oropharynx, can form abscesses, which may alter the spectrum of related clinical disease. The most common infections caused by the *S. anginosus* group are empyema, infective endocarditis, cellulitis, and abscesses in the head, neck, lung and abdomen (Kowlessar et al., 2006; Noguchi et al., 2015).

Many diseases are caused by or associated with streptococci, but the treatment

options are limited. Currently, one of the biggest problems is the bacterial antibiotic resistance. It has been reported that some GAS isolates are resistant to macrolides, tetracycline and fluoroquinolones (Walker et al., 2014). Among GBS, several isolates resulted resistant to clindamycin and macrolides, and the percentage may be as high as 30% in some areas (Compain et al., 2014). The predominant antibiotic resistance of *S. dysgalactiae* subsp. *equisimilis* isolates are erythromycin, clindamycin and tetracycline (Lu et al., 2016).

Some bacterial strains have intrinsic resistance, while others gain antibiotic resistance by mutation, horizontal gene acquisition and recombination with foreign DNA. In most instances, these mechanisms occur together. Mutations can be considered the workhorse of bacteria evolution, while the horizontal gene transfer (HGT) can immediately transfer a resistant gene (Perry et al., 2014). HGT is the sharing of genetic material among organisms other than by the DNA transmission from parent to offspring, which enable organisms to acquire new genes and phenotypes.

Microbe employs different mechanisms to transfer genes horizontally, including transformation, transduction and conjugation. In natural transformation, cells uptake naked DNA from the environment. It was reported that there exist about 80 species capable of transformation, and they are evenly divided between Gram-positive and Gram-negative bacteria (Johnston et al., 2014). Transduction is a process in which foreign DNA is transferred into a cell with the help of a virus or viral vector, transferring virulence genes between bacteria. Physical contact

between the cell providing the DNA and the cell receiving the DNA is not required. Conjugation is the transfer of DNA between cells by a bridge-like connection or by direct cell-to-cell contact. During conjugation the cell donates a conjugative or mobilizable genetic element that is often a transposon or a plasmid. The result is frequently beneficial to the recipient, gaining xenobiotic tolerance, antibiotic resistance or new metabolic capability. Agents responsible of DNA movement in HGT are called mobile genetic elements (MGEs). Here, we reviewed principal MGEs in streptococci species, including transposable elements, integrons, plasmids and bacteriophages.

Transposable elements

A transposable element, is a segment of DNA that can be integrated in different sites along a chromosome, sometimes creating or reversing mutations and altering the cell genetic identity and genome size. Transposable elements were discovered by the American geneticist McClintock in 1951 (Korona-Glowniak et al., 2015), and were widely recognized in 1967 when Shapiro found transposable elements in *Escherichia coli*. Initially transposable elements were considered unusual (Fedoroff, 2012a), but then they were regarded as 'selfish DNA' being able to replicate autonomously. More evidences showed that transposons play a crucial role in genome evolution (Fedoroff, 2012b). Transposable elements are commonly found in chromosomes, integrative conjugal elements (ICEs), bacteriophages, and plasmids. In prokaryotes, transposons can mainly fall into two categories, the insertion sequence (IS) and transposon (Tn).

Insertion sequences

An insertion sequence, also known as insertion sequence element, or IS, is a short DNA sequence that acts as a simple transposable element. It is defined as a small (2.5-kb), phenotypically cryptic segments of DNA with a simple genetic organization, and capable of inserting at multiple sites in a target molecule (Mahillon and Chandler, 1998). ISs present two major characteristics: they are small compared with other transposable elements and they only encode proteins relevant for transposition activity. These proteins usually catalyze enzymatic reactions, help the ISs movements, and can also stimulate or inhibit the transposition activity. However, in many cases, the exact role and activity of the protein are still unclear (Siguier et al., 2014).

Many studies have been conducted about the relationship between ISs and the antibiotic resistance in *Streptococcus*. Beigverdi et al. isolated forty-one *S. agalactiae* from pregnant women at 35-37 weeks of gestation, and found that 97.6%, 24.4%, and 14.6% of them were resistant to tetracycline, erythromycin and clindamycin, respectively. The insertion sequence IS1548 was found in 63.4% of isolates, which flanking the antibiotic gene (Beigverdi et al., 2014). Another study showed that the presence of rib, IS861 and GBSi1 group II intron within the C5a peptidase gene might accentuate the group B *Streptococcus* virulence potential and ecological success (Florindo et al., 2014). TnSs1 flanked by direct repeats of an IS6-family element may be related to the chloramphenicol resistance of *Streptococcus suis* (Takamatsu et al., 2003). In foodborne strains of *S. bovis*, there was a subregion of about 4 kb containing a novel mosaic tetracycline resistance gene *tet*(S/M) flanked by two copies of the IS1216 mobile

element (Barile et al., 2012).

Transposons

Transposons can be described as either composite transposons or complex transposons (Bennett, 2004). Composite transposons, for example Tn5, Tn9 and Tn10, are flanked by two separate identical or different IS elements. A composite transposon contains protein coding sequences encoding resistance to one or more antibiotics. A complex transposon has a more complicated structure than a composite transposon, being usually flanked by short IRs (Mahillon and Chandler, 1998). Between IRs there are genes related to transposition and antibiotic resistance (table.1).

Table 1. Details of Transposons discussed in this review

Name	Approximate size (kb)	Antibiotic resistance gene
Tn916	16.4	<i>tet</i> (M)
Tn917	5.3	<i>tet</i> (M) and <i>erm</i> (B)
Tn3872	24	<i>tet</i> (M) and <i>erm</i> (B)
Tn5801	25.8	<i>tet</i> (M)
Tn6002	20.9	<i>erm</i> (B) and <i>tet</i> (M)
Tn6003	25.1	<i>tet</i> (M) and <i>erm</i> (B)
Tn1545	25.3	<i>erm</i> (AM), <i>aphA</i> -3 and <i>tet</i> (M)
Tn2010	26.4	<i>mega</i> , <i>tet</i> (M) and <i>erm</i> (B)
Tn2009	23.5	<i>mega</i> , <i>tet</i> (M)
Tn1207.1	7.2	<i>mef</i> (A)
Tn1207.3	52.5	<i>mef</i> (A)
Tn2017	28.5	<i>mega</i> , <i>tet</i> (M), <i>erm</i> (B)
Tn5251	18	<i>tet</i> (M)
Tn5253	64.5	<i>cat</i> and <i>tet</i> (M)
Tn5252	47.5	<i>cat</i> and <i>tet</i> (M)

Tn916

Tn916 is a 16.4 kb transposon containing a tetracycline resistance determinant. Tn916 can transfer through conjugation at frequencies of 10^{-8} to 10^{-5} per donor in the absence of plasmid DNA, and it is also capable of insertion at different locations of the recipient chromosome (Johnson and Grossman, 2015). Some of the trans-conjugants possess more than two non-tandem copies (Gawron-Burke and Clewell, 1982), and have been named “conjugative transposon”, or “integrative conjugative element”. The recombination module of all Tn916-related elements is located at one end of the element with imperfect repeat sequences. Tn916 contains four functional modules: regulation, conjugation, recombination and accessory genes not involved in conjugative transposition (e.g. tetracycline resistance). The proposed regulatory region of Tn916 includes *orf12*, *orf9*, *orf7* and *orf8*. *Orf6* and *orf10* are also in this region, but their functions are not clear yet (Figure.1) (Celli and Trieu-Cuot, 1998). Homologous segments have been found in the transposon junction regions but there was no direct duplication of the target sequence (Clewell et al., 1988). Tn916 replicates autonomously *via* a rolling-circle mechanism, which depends on the relaxase encoded by *orf20* (Wright and Grossman, 2016). It has been reported that Tn916 transposes by an excision-insertion mechanism (Caparon and Scott, 1989), in which it is firstly excised by a cut from donor cells and transformed into a circular form, then it is transferred by cell-cell contact, and finally it is inserted into receiving cells through recombinase (Roberts and Mullany, 2011). Two transposon-encoded proteins, Xis-Tn (an accessory protein) and Int-Tn (a tyrosine recombinase), are needed for excision. Int-Tn alone is enough for integration. The conjugative transposition of Tn916 needs a functional integrase in both recipient and donor strain (Storrs et

al., 1991). The DNA sequences of the target sites of many Tn916 insertions showed that Tn916 prefers to use A-T rich targets in different hosts (Wang et al., 2000). The most important regulatory event in Tn916 for conjugative transposition is the interaction between ribosomes and tetracycline, which makes the transcriptional terminator destroyed at the end of *orf12* and continues transcribing the *tet(M)* and the downstream genes. Tetracycline can inactivate most of the ribosomes, and a few ribosomes can remain active by the protection of the basal expression of *tet(M)* (Roberts and Mullany, 2009).




Figure 1. Schematic representation of Tn916.

The Tn916 family is widespread among numerous commensal and pathogenic bacteria. Tn916 has an extensive host range, both *in vitro* and *in vivo* (Scott, 2002). In *S. suis* serotype 2, the presence of Tn916 with *tet(M)* has favored the pathogenicity and dissemination of this pathogen (Zhang et al., 2015). In 138 antibiotic resistant pneumococcal isolated strains from nasopharynx of healthy young children from Lublin of Poland, Tn916 was detected in 16 strains (Korona-Glowniak et al., 2015). Moreover, among 62 tetracycline-resistant *tet(M)*-positive *S. agalactiae* isolate strains, 70% of them harboring Tn916 (Mingoia et al., 2013).

Tn917

Tn917 was identified on pAD2, a nonconjugative plasmid of *Enterococcus faecalis* strain DS16 (Tomich et al., 1980). Erythromycin resistance associated with Tn917 is inducible, and the exposition of cells to low concentrations of erythromycin can induce transposition from pAD2 to pAD1. Tn917 is 5,257 bp in length and has six open reading frames. *Orf2* is *erm(B)*, which is responsible for erythromycin resistance. In 263 viridans group streptococci strains, the long-established transposon Tn917 has been found in seven isolates by PCR mapping, which including one *Streptococcus mitis* strain, three *Streptococcus oralis* strains, two *S. sanguinis* strains and one *S. anginosus* strain (Brenciani et al., 2014).

Tn3872

Tn3872 is found by DNA-DNA hybridization, and nucleotide sequence analyses revealed that they were made up by the insertion of Tn917 into *orf9* of Tn916 (McDougal et al., 1998). With the *tet(M)* gene of Tn916 linked to the *erm(B)* gene of Tn917, Tn3872 can lead to the erythromycin and tetracycline resistance. *Tet(M)* on the Tn3872-like transposon in the *S. agalactiae* isolates has been confirmed by PCR with the primers *tet(M)*-forward and *erm(B)* (Sadowy et al., 2010). Thirteen strains had Tn3872 among twenty-one clinical *Streptococcus salivarius*, and two of them carried a Tn3872 with a silent *tet(M)* gene (Chaffanel et al., 2015). In pneumococcus, *tet(M)* can evolve through homologous recombination after Tn3872 integration into the host chromosome (Chaffanel et al., 2015). It has been reported that Tn916 can transfer from an enterococcal donor to a pneumococcal recipient, but the composite transposon Tn3872 cannot transfer from a pneumococcal to another pneumococcal or from an enterococcal to a pneumococcal. The low transfer frequency of Tn3872 or the inactivation of

mobility of Tn916 due to the insertion of Tn917 may be the reason (Swartley et al., 1993). However, in *Streptococcus defectives* strains, the frequency of conjugative transposition of Tn3872 was ranging from 4×10^{-6} to 2.8×10^{-8} (Poyart et al., 2000). Tn3872-related elements were distributed in many serotypes of pneumococcus, i.e. 6B, 23F, and 14 (Quintero et al., 2011; Ramos et al., 2014). In some strains, Tn3872 and Tn3872-related elements were lacking *tet(M)* gene (Cochetti et al., 2007; Quintero et al., 2011). Genetic elements relevant to drug resistance can insert in Tn3872, and the strain will display resistance to other antibiotics. For example, *mega*, a genetic element carrying *mef(E)*, has been found transferred by conjugation to *S. pneumoniae* and then linked with Tn3872-like transposon by recombinases (Figure.2) (Del Grosso et al., 2006; Santagati et al., 2009). In some strains, Tn3872 carries transposase genes *tnpA* and *tnpR* (Korona-Glowniak et al., 2016; Korona-Glowniak et al., 2015).




Figure 2. *Mega* of *S. pneumoniae*.

Tn5801

Tn5801, 25.8 kb in size, was originally detected in *Staphylococcus aureus* Mu50 (Kuroda et al., 2001). It is a Tn916 family element, and the *int* and *xis* genes of Tn916 are replaced by a unique gene (*int5801*) encoding different types of tyrosine recombinases. The *tet(M)* genes of Tn5801 and Tn916 are similar, with

about 97.7% DNA identity (de Vries et al., 2016). Tn5801-like transposons have been detected in some streptococci, including *S. mitis* B6 (Denapaité et al., 2010), *S. agalactiae* (Da Cunha et al., 2014; Mingoia et al., 2013) and *S. oralis* (Brenciani et al., 2014). In *tet*(M)-positive tetracycline resistant *S. agalactiae*, 30% harbors Tn5801-like element (Mingoia et al., 2013).

Tn6002

Tn6002, a 20.9 kb element, is formed by the insertion of an *erm*(B)-carrying DNA fragment into Tn916, which is located between *orf20* and *orf19* (Brenciani et al., 2007). It was originally discovered in *Streptococcus cristatus* and designated as Tn916Erm, whose complete sequence is available in GenBank since 2005 (accession no. AY898750) (Varaldo et al., 2009). It has also been found in *S. pyogenes* (Ardanuy et al., 2010; Brenciani et al., 2007), *S. salivarius* (Chaffanel et al., 2015) and *S. pneumoniae* (Calatayud et al., 2007; Korona-Glowniak et al., 2016; Quintero et al., 2011). In *erm*(B)-positive pneumococci, Tn6002 was the most common *erm*(B)-carrying Tn916-related element (Cochetti et al., 2008; Korona-Glowniak et al., 2015). The spread of Tn6002 from the ST386/ST4310/ST4825 lineage to the serotype 6C of the pneumococci isolates lead to the increase of the antimicrobial non-susceptibility of serotype 6C in the late-PCV7 period (Rolo et al., 2011). In viridans group streptococci (VGS), Tn6002 were also detected into *S. mitis* and *S. oralis* (Brenciani et al., 2014). In oral streptococci isolated from healthy human, Tn6002 was detected from *Streptococcus infantis*, *S. mitis*, *S. sanguinis*, *Streptococcus australis*, *S. oralis* and *Streptococcus parasanguini*.

Tn6003

Tn6003, a new Tn6002-related composite element, found in 2007 (Cochetti et al., 2007), contains *tet(M)* and *erm(B)* gene, as well as the *aadE-sat4-aphA-3* gene cluster that forms a 4.2 kb fragment called MAS (macrolideaminoglycoside-streptothricin). Tn6003 is 25.1 kb (accession number: AM410044). The transfer frequency of Tn6003 from *S. pneumoniae* strain Ar4 to *E. faecalis* JH2-2 is 1.7×10^{-7} (Cochetti et al., 2007). Tn6003 is a common Tn916-related element in *erm(B)*-positive pneumococci (Quintero et al., 2011).

Tn1545

Tn1545, a 25.3 kb element, was originally detected in the chromosome of *S. pneumoniae* BM4200 (an antibiotics-resistant clinical isolates) (Courvalin and Carrier, 1987). It confers resistance to erythromycin (*ermAM*), kanamycin (*aphA-3*) and tetracycline (*tetM*) and was the first genetic element found carrying the *erm(B)* gene in *S. pneumoniae*. Compared with Tn916, between *orf20* and *orf19*, Tn1545 showed the insertion of two *erm(B)* and MAS elements; and between *orf13* and *tet(M)*, Tn1545 had the insertion of IS1239 (Quintero et al., 2011). Tn1545 is self-transferable to various streptococci, including *Streptococcus cremoris*, *S. sanguinis*, *Streptococcus lactis*, *Streptococcus diacetylactis*, and *E. faecalis*. It is a conjugative shuttle transposon, which can transpose from the hybrid plasmid to various new sites of the new host chromosome and then re-transpose from the chromosome to the plasmid at a frequency of 5×10^{-9} . Though the absence of recombinase genes, the MAS element of Tn1545 is able to exploit

long flanking repeats for excision in a circular form (Palmieri et al., 2012).

Tn2010

Tn2010, 26.4 kb in length, is a composite element carrying both *mega*, *tet(M)* and *erm(B)* genes (Del Grosso et al., 2007; Talebi et al., 2016). Tn2010 is similar to Tn6002, including sequence and structure, and it has the insertion of the *mega* element into *orf6* of the Tn6002, as well as 70 nucleotide mutations (Li et al., 2011). It has been indicated that Tn2010 could not transfer macrolide resistance through conjugation (Del Grosso et al., 2006). Zhou et al. proved that Tn2010 is more likely to be transferred by transformation other than conjugation in *S. pneumoniae*, and the transformation frequency is 3×10^{-7} transformants per cfu (Zhou et al., 2014).

Tn2010 is common in *S. pneumoniae* (Bowers et al., 2012; Korona-Glowniak et al., 2015), and recently it has been found in *S. mitis* (Brenciani et al., 2014). In *S. pneumoniae*, most strains owning Tn2010 belonging to serotype 19A or 19F (Korona-Glowniak et al., 2016). The wide spreading of the macrolide resistance genes is highly relevant to Tn2010. *S. pneumoniae* A026, isolated from cerebrospinal fluid, contains Tn2010 and is resistant to penicillin, trimethoprim, erythromycin, sulfonamides, and tetracycline but is susceptible to levofloxacin (Sui et al., 2013).

Tn2009

Tn2009 is a composite element of approximately 23.5 kb, originally found in the chromosomes of the two *S. pneumoniae* strains (Del Grosso et al., 2004). Subsequently, it was detected in *S. mitis* (Brenciani et al., 2014), *S. pyogenes* (Del Grosso et al., 2011) and *S. salivarius* (Chaffanel et al., 2015). Tn2009 has the insertion of the *mega* element into the *orf6* of a Tn916-like transposon containing *tet(M)*. In many Tn2009 element of *S. pyogenes* strains, *mef(E)* is included in a typical *mega* element, while in some strains, *mef(E)* is physically linked with *tet(M)* (Del Grosso et al., 2011). Tn2009 contains determinants for tetracycline, macrolide and erythromycin resistance. In *S. pneumoniae* and *S. pyogenes* the conjugal transfer of Tn2009 has not been detected, unlike the transfer by transformation (Del Grosso et al., 2016; Del Grosso et al., 2004). It has been reported that the conjugative Tn2009 in a Gram-negative commensal bacteria *Acinetobacter junii* 326 was indistinguishable from the *S. pneumoniae* Tn2009 element (Ojo et al., 2006).

Tn1207.1

Tn1207.1, 7,244 bp in size, is the first described *mef*-carrying genetic element. It has been found in both *S. pneumoniae* and *S. pyogenes* (Santagati et al., 2000). Tn1207.1 consists in first 5 ORFs with the same direction of transcription and the last 3 ORFs in the opposite direction (Pozzi et al., 2004). The *orf2* encodes a site-specific recombinase, while *orf5*, starting from the downstream of the *mef(A)*, is homologous to *vga(A)* and *msr(SA)* (Pozzi et al., 2004). The *mrs(SA)* and *mef(A)* are two important parts of the *mega* element. Tn1207.1 is not a conjugative element (Santagati et al., 2000).

Tn1207.3

Tn1207.3, homologous to Tn1207.1, is resistant to macrolides and carries the *mef(A)* gene. Tn1207.3 was found in *S. pyogenes*, *S. pneumoniae*, *S. dysgalactiae* subsp. *Equisimilis* (Figueiredo et al., 2006; Marimon et al., 2005; Rato et al., 2010), *S. gordonii* (Santagati et al., 2003) and *S. agalactiae* (Figueiredo et al., 2006; Marimon et al., 2005). Its nucleotide sequence is highly identical to that of Tn1207.1, and only the right end of the Tn1207.3 has a new sequence (Pozzi et al., 2004; Santagati et al., 2003). Tn1207.3 can transfer from *S. pyogenes* to *S. pneumoniae* by conjugation, and from *S. pneumoniae* trans-conjugant MF2 to *S. pyogenes* and *S. gordonii* (Santagati et al., 2003). The transfer frequency of Tn1207.3 to various *S. pyogenes* is significantly different, and the mean frequencies range from 7.20×10^{-8} to 1.13×10^{-6} conjugant per recipient (Hadjirin et al., 2014). The integration of Tn1207.3 did not cause deletions, while integration of Tn1207.1 caused a 1947 bp deletion (Pozzi et al., 2004). The DNA sequencing of Tn1207.3 indicated that it is actually a prophage, the same with the right end of $\phi 10394.4$ (Iannelli et al., 2014a).

Tn2017

Tn2017, a Tn916-like element, was found in *S. pneumoniae* and beta-hemolytic *Streptococcus* (Korona-Glowniak et al., 2015). Tn2017 resembles Tn3872, which is the insertion of *mega* element into the *orf6* of Tn3872, thus being resistant to erythromycin, tetracycline and macrolide. Tn2017 also contains the *tnpA* and *tnpR* genes. In erythromycin and tetracycline resistant beta-hemolytic

Streptococcus isolates, Tn2017 is the most prevalent transposon, while in *S. pneumonia* isolates it is the second most common transposon (Tantivitayakul et al., 2016).

Tn5251

Tn5251, 18 kb in length, is like Tn916 in structure and size. It contains 22 ORFs, of which 20 have the same direction and transcription, and half are annotated (Santoro et al., 2010). Tn5251 carries *tet(M)*, *int* and *six* genes. Tn5251 cannot transfer independently from Tn5253 through conjugation (Kilic et al., 1994), however recently it has been proposed that Tn5251 can transfer autonomously in *S. pneumoniae* and *E. faecalis*, and the autonomous copies can be moved into *S. pneumoniae*, *S. gordonii*, *S. agalactiae*, *S. pyogenes*, *E. faecalis* and *Bacillus subtilis* (Santoro et al., 2010).

Tn5253

Tn5253, originally found in the chromosome of *S. pneumoniae* BM6001 (Dang-Van et al., 1978), was formerly called the Ω (*cat-tet*) element, carrying the *cat* and *tet(M)* genes (Vijayakumar et al., 1986). Tn5253 is 64.5 kb in size, and contains 79 ORFs in total, 38 of which have annotations (Iannelli et al., 2014b). Tn5251 and Tn5252 were found integrated into the Tn5253, demonstrating that Tn5253 was a composite element (Ayoubi et al., 1991). The highest conjugation frequency of Tn5253 was 6.7×10^{-3} trans-conjugants/donor when *S. pyogenes* was the donor (Iannelli et al., 2014b). Tn5253 is common in the drug resistance of *S. pneumoniae* strains, especially in the serotype 6B strains, but the *cat* gene may

not be detected for the modifications of the transposon (Quintero et al., 2011). Among clinical isolates of *S. pneumoniae*, Tn5253-related elements are extremely different (Henderson-Begg et al., 2009; Mingoia et al., 2011). In streptococci, *cat(Q)* and *mef(I)* are linked in a genetic module called IQ module that has been detected in *S. pneumoniae*, *S. pyogenes* and viridans group streptococci. The IQ module in *S. pyogenes* belongs to the Tn5253 family (Mingoia et al., 2014a; Mingoia et al., 2014b; Mingoia et al., 2007).

Tn5252

Tn5252, 47.5 kb in size, is a streptococcal conjugative transposon (Kilic et al., 1994). It was originally found in the chromosome of *S. pneumoniae* BM6001 as a part of Tn5253 (Shoemaker et al., 1979). Until now, Tn5252 has only been partially sequenced, showing the presence of an integrase gene, *tet(M)*, and *cat*. While Tn916 class of element can insert at several spots, Tn5252 prefer to integrate at a specific site, which is drove by *int* gene. *Cat* gene can encode chloramphenicol acetyltransferases (CATs) that lead to the enzymatic inactivation of the drug, and Tn5252 can be detected in the clinical chloramphenicol resistant strains (Korona-Glowniak et al., 2015). Since the discovery of Tn5252, its homologues have been detected in many streptococcal species, such as *S. suis*, *S. pneumoniae* and GBS (Kayansamruaj et al., 2015; Wyres et al., 2013).

Integron

Integrans are genetic elements that can help bacteria to enhance their ability of adaptation and evolution through the acquisition, stockpiling, excision and

differential expression of new genes (Escudero et al., 2015). The genes are usually contained in a gene structure called cassette, a circular element carrying one promoterless gene together with a recombination site (*attC*) (Figure.3). Another structure included in integrons is a stable platform. The platform has several parts, and the first is *intI*, a gene encoding the integrase, which is a member of tyrosine recombinase family (Messier and Roy, 2001). The second part is the *attI* site, an integron-associated recombination site, which is in the upstream of the *intI* gene (Partridge et al., 2000). The third part is *Pc*, an integron-associated promoter, located between the *intI* and the *attI* site or within the *intI* gene (Collis and Hall, 1995). Cassettes are non-replicative elements, and they play their roles by integrating into the platform and expression from the *Pc* promoter. With the constant integration taking place, an array of cassettes will be assembled, and the cassettes' excised and integrated into the first position have the maximal expression (Escudero et al., 2015). There are two key advantages owning by the integron system for the genomic studies. First, the new cassette is integrated into the bacterial genome at *attI* site, which will not disturb the existing genes. Second, the newly integrated cassettes are expressed by the promoter *Pc* that it is ready for the natural selection (Gillings, 2014).

At present, there are five classes of mobile integrons associated with antibiotic resistance: classes 1, 2 and 3, obtained from clinical contexts (Partridge, 2011); class 4, detected in *Vibrio cholerae*; and class 5, found on the plasmid of *Alivibrio salmonicida* (Gillings, 2014). All five classes are linked to mobile elements, such as conjugative plasmids, insertion sequences and transposons, which can work as vehicles for the gene transmission (Mazel, 2006).

Integrons play an important role in the spread of antibiotic resistance, especially

in Gram-negative pathogens. Nonetheless, recent studies focused on the role of integrons in antibiotic resistance in Gram-positive microorganisms (Deng et al., 2015; Li et al., 2017), such as *Staphylococcus*, *Enterococcus*, *Corynebacterium* and *Streptococcus*. In streptococci, *S. pneumoniae* was proved to carry class 1 integron (Shi et al., 2006).




Figure 3. General structure of class 1 integrons.

Plasmids

Plasmids are a group of extrachromosomal DNA molecules that can self-replicate, are usually circular and does not contain genes required for essential cellular functions. The typical plasmid is a circular, closed double-stranded DNA molecule; nonetheless, linear double-stranded plasmids have been found (Stewart et al., 2005). In general, a plasmid includes essential genes related to replicative functions and accessory genes encoding different functions (Fernandez-Lopez et al., 2006) including drug resistance, fitness and virulence factors (Gyles and Boerlin, 2014). Plasmids are inherited from parent to daughter cells at cell division by partitioning. For some plasmids carrying accessory genes, they are therefore large and always present in low copy numbers in their host to ensure that the metabolic burden of maintaining and duplicating their genomes are not excessive. And the low copy number would increase their chance of being faithfully transmitted to daughter cells during division. Therefore, in order to decrease

random losses, the plasmid should be inherited stably at a short time (Sengupta and Austin, 2011).

Some plasmids carry genes allowing them to transfer by conjugation, while others are lacking such genes and count on the transferable function of other plasmids (Smillie et al., 2010). The conjugative genes trigger DNA transfer from the donor cell to the recipient cell, and other genes guarantee the DNA survival in the new host. The transfer genes also encode proteins able to 'pump' the DNA into recipient cells, and these proteins all belong to the TraG-like family of ATPases (Schroder et al., 2002). Relaxase are also important in many conjugative systems that can nick DNA to give a single-stranded substrate suitable for transfer, and the nicking process takes place in a specific site. In Gram-negative bacteria the pilus is an important section of conjugative systems. Pilus assembly plays the role of a type IV secretion system (T4SS) (Lybarger and Sandkvist, 2004), which help bacteria in two aspects: the genetic exchange and the transmission of effector molecules to target cells (Cascales and Christie, 2003).

In Gram-positive bacteria conjugative mechanisms include the pheromone-independent system that have a cell surface protein able to trigger mating-pair formation and pheromone-dependent mating systems in enterococci (Frost et al., 2005). In streptococci, the transfer of chromosomal DNA may be affected by the plasmid transfer in horizontal gene transfer, due to the chromosome mobilization ability of plasmids and the high frequency of recombination mode of the F plasmids (Lawley et al., 2003). Plasmids have been grouped according to their

replicons, and the streptococcal plasmid pMV158 belongs to the Rep_2 family (PF01719). There are several parts related to the mobilization of plasmid pMV158: prototype of the MOBV relaxase family, the product of the *mobM* gene, and its cognate origin of transfer, *oriT*. The *mobM* gene encodes the relaxase MobM, the relaxase of pMV158. MobM forms a dimer through its C-terminal domain, anchoring the protein to the cell membrane and participating in T4SS protein-protein interactions (Fernandez-Lopez et al., 2013). Plasmid pLS1 is a deleted derivative of pMV158, which has lost the ability of conjugative mobilization, thus decreasing the host cell growth by 8-9% in *S. pneumonia* (Hernandez-Arriaga et al., 2012). Many toxins and drug resistance genes in streptococci are encoded by plasmid. The β plasmid from *E. faecalis* can transfer the erythromycin and lincomycin resistance genes by conjugation (LeBlanc et al., 1978). The diplococci-producing ability can be transferred from the plasmid of *Streptococcus cremoris* 346 to the plasmid-free *S. cremoris* by conjugation with high frequency (Davey, 1984). The minocycline resistance in an oral *Streptococcus infantis* isolate may be due to the *tet(S)* gene which is carried by a novel low copy number plasmid (Ciric et al., 2014).

Bacteriophages

Certain drug resistance in bacteria are found to be relevant to phages (Brussow et al., 2004; Lekunberri et al., 2017), and there are many antibiotic resistant genes carried by the bacteriophages, such as *tetA*, *tetW*, *intl1*, *intl2*, *intl3*, *blaOXA-2* and *blaTEM* (Anand et al., 2016). Phage genomes may contain DNA or RNA, and the DNA can be single- or double-stranded. The essential genes of phages include the replicase genes, genes encoding for proteins involved in the DNA packaging,

and genes encoding phage components used to disturb the host cell replicative machinery (Frost et al., 2005).

Phage-mediated horizontal gene transfer occurs via lysogenic conversion or transduction. Moreover, some phages can play a significant role in the emergence of pathogens, including *S. aureus* (food poisoning), *S. pyogenes* (scarlet fever), *E. coli* (Shiga toxin) and *Clostridium botulinum* (botulism). In *S. mitis*, phage tail genes *pbIA* and *pbIB* encode adhesion proteins for bacterial host attachment. The process of phage DNA packaging is highly accurate, but the host genome DNA fragment may be packaged at a finite frequency compared with the phage DNA, resulting in phage particles that can attach and deliver the DNA into suitable bacteria, with the injected exogenous bacterial DNA integrated into the genome. The observation of toxin genes in *S. pyogenes* and *Corynebacterium* located next to the phage attachment site raised the hypothesis that the prophage genes actually stand for bacterial genes obtained by a false excision from a previous bacterial host (Brussow et al., 2004).

There are likely many new variants of MGEs, which can contain other resistant genes able to transfer to broad host range. Besides, MGEs may evolve complicated mechanisms of transcriptional regulation which likely minimize the biological cost to the host. Furthermore, many of the current studies are focus on laboratory strains, which may not reflect the variability of natural isolates. Therefore, it is urgent to clarify the structure and function of the DNA integrated in a new host, the rate of the gene transfer in both laboratory and natural bacteria,

as well as the effects of MGEs on the transcriptional regulation of the gene expression.

Conclusions

Here, we have reviewed principal MGEs in streptococci, including transposable elements, integrons, plasmids, and bacteriophages. MGEs have a strong environmental and clinical impact due to their role in the spread of antibiotic resistance in different bacteria. Transposable elements, mainly containing IS and transposons, are strictly related to the resistance to erythromycin, tetracycline, kanamycin and macrolide. The relationship between antibiotic resistance and integrons is well studied. Recently class 1 integron has been found in *S. pneumoniae*. Many toxins and the spread of drug resistance in *Streptococcus* are plasmid encoded, including the resistance to erythromycin, minocycline, and lincomycin. Finally, phages can mediate horizontal gene transfer *via* lysogenic conversion or transduction.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (31200985 and 31870065). The authors declare no potential conflicts of interest regarding the authorship and/or publication of this article.

References

Anand, T., Bera, B.C., Vaid, R.K., Barua, S., Riyesh, T., Virmani, N., Hussain, M., Singh, R.K., and Tripathi, B.N. (2016). Abundance of antibiotic resistance genes in environmental bacteriophages. *J Gen Virol* 97, 3458-3466 DOI: 10.1099/jgv.0.000639.

Ardanuy, C., Domenech, A., Rolo, D., Calatayud, L., Tubau, F., Ayats, J., Martin, R., and Linares, J. (2010). Molecular characterization of macrolide- and multidrug-resistant *Streptococcus pyogenes* isolated from adult patients in Barcelona, Spain (1993-2008). *J Antimicrob Chemother* 65, 634-643 DOI: 10.1093/jac/dkq006.

Ayoubi, P., Kilic, A.O., and Vijayakumar, M.N. (1991). Tn5253, the pneumococcal omega (*cat tet*) BM6001 element, is a composite structure of two conjugative transposons, Tn5251 and Tn5252. *J Bacteriol* 173, 1617-1622

Barile, S., Devirgiliis, C., and Perozzi, G. (2012). Molecular characterization of a novel mosaic *tet*(S/M) gene encoding tetracycline resistance in foodborne strains of *Streptococcus bovis*. *Microbiology* 158, 2353-2362 DOI: 10.1099/mic.0.058206-0.

Beigverdi, R., Jabalameli, F., Mirsalehian, A., Hantoushzadeh, S., Boroumandi, S., Taherikalani, M., and Emaneini, M. (2014). Virulence factors, antimicrobial susceptibility and molecular characterization of *Streptococcus agalactiae* isolated from pregnant women. *Acta Microbiol Immunol Hung* 61, 425-434 DOI: 10.1556/AMicr.61.2014.4.4.

Bennett, P.M. (2004). Genome plasticity: insertion sequence elements, transposons and integrons, and DNA rearrangement. *Methods Mol Biol* 266, 71-113 DOI: 10.1385/1-59259-763-7:071.

Bowers, J.R., Driebe, E.M., Nibecker, J.L., Wojack, B.R., Sarovich, D.S., Wong, A.H., Brzoska, P.M., Hubert, N., Knadler, A., Watson, L.M., et al. (2012). Dominance of multidrug resistant CC271 clones in macrolide-resistant *streptococcus pneumoniae* in Arizona. BMC Microbiol 12, 12 DOI: 10.1186/1471-2180-12-12.

Brandt, C.M., and Spellerberg, B. (2009). Human infections due to *Streptococcus dysgalactiae* subspecies *equisimilis*. Clin Infect Dis 49, 766-772 DOI: 10.1086/605085.

Brenciani, A., Bacciaglia, A., Vecchi, M., Vitali, L.A., Varaldo, P.E., and Giovanetti, E. (2007). Genetic elements carrying *erm(B)* in *Streptococcus pyogenes* and association with *tet(M)* tetracycline resistance gene. Antimicrob Agents Chemother 51, 1209-1216 DOI: 10.1128/AAC.01484-06.

Brenciani, A., Tiberi, E., Tili, E., Mingoia, M., Palmieri, C., Varaldo, P.E., and Giovanetti, E. (2014). Genetic determinants and elements associated with antibiotic resistance in viridans group streptococci. J Antimicrob Chemother 69, 1197-1204 DOI: 10.1093/jac/dkt495.

Brussow, H., Canchaya, C., and Hardt, W.D. (2004). Phages and the evolution of bacterial pathogens: from genomic rearrangements to lysogenic conversion. Microbiol Mol Biol Rev 68, 560-602, table of contents DOI: 10.1128/MMBR.68.3.560-602.2004.

Calatayud, L., Ardanuy, C., Cercenado, E., Fenoll, A., Bouza, E., Pallares, R., Martin, R., and Linares, J. (2007). Serotypes, Clones, and Mechanisms of Resistance of Erythromycin-Resistant *Streptococcus pneumoniae* Isolates Collected in Spain. Antimicrob Agents Chemother 51, 3240-3246 DOI:

10.1128/AAC.00157-07.

Caparon, M.G., and Scott, J.R. (1989). Excision and insertion of the conjugative transposon Tn916 involves a novel recombination mechanism. *Cell* 59, 1027-1034

Cascales, E., and Christie, P.J. (2003). The versatile bacterial type IV secretion systems. *Nat Rev Microbiol* 1, 137-149 DOI: 10.1038/nrmicro753.

Celli, J., and Trieu-Cuot, P. (1998). Circularization of Tn916 is required for expression of the transposon-encoded transfer functions: characterization of long tetracycline-inducible transcripts reading through the attachment site. *Mol Microbiol* 28, 103-117

Chaffanel, F., Charron-Bourgoin, F., Libante, V., Leblond-Bourget, N., and Payot, S. (2015). Resistance genes and genetic elements associated with antibiotic resistance in clinical and commensal isolates of *Streptococcus salivarius*. *Appl Environ Microbiol* 81, 4155-4163 DOI: 10.1128/AEM.00415-15.

Ciric, L., Brouwer, M.S., Mullany, P., and Roberts, A.P. (2014). Minocycline resistance in an oral *Streptococcus infantis* isolate is encoded by *tet(S)* on a novel small, low copy number plasmid. *FEMS Microbiol Lett* 353, 106-115 DOI: 10.1111/1574-6968.12410.

Claridge, J.E., 3rd, Attorri, S., Musher, D.M., Hebert, J., and Dunbar, S. (2001). *Streptococcus intermedius*, *Streptococcus constellatus*, and *Streptococcus anginosus* ("Streptococcus milleri group") are of different clinical importance and are not equally associated with abscess. *Clin Infect Dis* 32, 1511-1515 DOI: 10.1086/320163.

Clewell, D.B., Flannagan, S.E., Ike, Y., Jones, J.M., and Gawron-Burke, C. (1988). Sequence analysis of termini of conjugative transposon Tn916. *J Bacteriol* 170, 3046-3052

Cochetti, I., Tili, E., Mingoia, M., Varaldo, P.E., and Montanari, M.P. (2008). *erm(B)*-carrying elements in tetracycline-resistant pneumococci and correspondence between Tn1545 and Tn6003. *Antimicrob Agents Chemother* 52, 1285-1290 DOI: 10.1128/AAC.01457-07.

Cochetti, I., Tili, E., Vecchi, M., Manzin, A., Mingoia, M., Varaldo, P.E., and Montanari, M.P. (2007). New Tn916-related elements causing *erm(B)*-mediated erythromycin resistance in tetracycline-susceptible pneumococci. *J Antimicrob Chemother* 60, 127-131 DOI: 10.1093/jac/dkm120.

Collis, C.M., and Hall, R.M. (1995). Expression of antibiotic resistance genes in the integrated cassettes of integrons. *Antimicrob Agents Chemother* 39, 155-162

Compain, F., Hays, C., Touak, G., Dmytruk, N., Trieu-Cuot, P., Joubrel, C., and Poyart, C. (2014). Molecular characterization of *Streptococcus agalactiae* isolates harboring small *erm(T)*-carrying plasmids. *Antimicrob Agents Chemother* 58, 6928-6930 DOI: 10.1128/AAC.03855-14.

Courvalin, P., and Carlier, C. (1987). Tn1545: a conjugative shuttle transposon. *Mol Gen Genet* 206, 259-264

Da Cunha, V., Davies, M.R., Douarre, P.E., Rosinski-Chupin, I., Margarit, I., Spinali, S., Perkins, T., Lechat, P., Dmytruk, N., Sauvage, E., et al. (2014). *Streptococcus agalactiae* clones infecting humans were selected and fixed through the extensive use of tetracycline. *Nat Commun* 5, 4544 DOI: 10.1038/ncomms5544.

Dang-Van, A., Tiraby, G., Acar, J.F., Shaw, W.V., and Bouanchaud, D.H. (1978). Chloramphenicol resistance in *Streptococcus pneumoniae*: enzymatic acetylation and possible plasmid linkage. *Antimicrob Agents Chemother* 13, 577-583

Davey, G.P. (1984). Plasmid associated with diplococcin production in *Streptococcus cremoris*. *Appl Environ Microbiol* 48, 895-896

de Vries, L.E., Hasman, H., Jurado Rabadan, S., and Agerso, Y. (2016). Sequence-Based Characterization of Tn5801-Like Genomic Islands in Tetracycline-Resistant *Staphylococcus pseudintermedius* and Other Gram-positive Bacteria from Humans and Animals. *Front Microbiol* 7, 576 DOI: 10.3389/fmicb.2016.00576.

Del Grosso, M., Camilli, R., Barbabella, G., Blackman Northwood, J., Farrell, D.J., and Pantosti, A. (2011). Genetic resistance elements carrying *mef* subclasses other than *mef(A)* in *Streptococcus pyogenes*. *Antimicrob Agents Chemother* 55, 3226-3230 DOI: 10.1128/AAC.01713-10.

Del Grosso, M., Camilli, R., Iannelli, F., Pozzi, G., and Pantosti, A. (2006). The *mef(E)*-carrying genetic element (mega) of *Streptococcus pneumoniae*: insertion sites and association with other genetic elements. *Antimicrob Agents Chemother* 50, 3361-3366 DOI: 10.1128/AAC.00277-06.

Del Grosso, M., Camilli, R., Rizzi, E., Pietrelli, A., De Bellis, G., and Pantosti, A. (2016). ICESpy009, a Conjugative Genetic Element Carrying *mef(E)* in *Streptococcus pyogenes*. *Antimicrob Agents Chemother* 60, 3906-3912 DOI: 10.1128/AAC.03082-15.

Del Grosso, M., Northwood, J.G., Farrell, D.J., and Pantosti, A. (2007). The macrolide resistance genes *erm(B)* and *mef(E)* are carried by Tn2010 in dual-

gene *Streptococcus pneumoniae* isolates belonging to clonal complex CC271. Antimicrob Agents Chemother 51, 4184-4186 DOI: 10.1128/AAC.00598-07.

Del Grosso, M., Scotto d'Abusco, A., Iannelli, F., Pozzi, G., and Pantosti, A. (2004). Tn2009, a Tn916-like element containing *mef(E)* in *Streptococcus pneumoniae*. Antimicrob Agents Chemother 48, 2037-2042 DOI: 10.1128/AAC.48.6.2037-2042.2004.

Denapaite, D., Bruckner, R., Nuhn, M., Reichmann, P., Henrich, B., Maurer, P., Schahle, Y., Selbmann, P., Zimmermann, W., Wambutt, R., et al. (2010). The genome of *Streptococcus mitis* B6--what is a commensal? PLoS One 5, e9426 DOI: 10.1371/journal.pone.0009426.

Deng, Y., Bao, X., Ji, L., Chen, L., Liu, J., Miao, J., Chen, D., Bian, H., Li, Y., and Yu, G. (2015). Resistance integrons: class 1, 2 and 3 integrons. Ann Clin Microbiol Antimicrob 14, 45 DOI: 10.1186/s12941-015-0100-6.

Escudero, J.A., Loot, C., Nivina, A., and Mazel, D. (2015). The integron: adaptation on demand. Microbiol Spectr 3, MDNA3-0019-2014 DOI:10.1128/microbiolspec.MDNA3-0019-2014.

Facklam, R. (2002). What happened to the streptococci: overview of taxonomic and nomenclature changes. Clin Microbiol Rev 15, 613-630

Fedoroff, N.V. (2012a). McClintock's challenge in the 21st century. Proc Natl Acad Sci U S A 109, 20200-20203 DOI: 10.1073/pnas.1215482109.

Fedoroff, N.V. (2012b). Presidential address. Transposable elements, epigenetics, and genome evolution. Science 338, 758-767

Fernandez-Lopez, C., Pluta, R., Perez-Luque, R., Rodriguez-Gonzalez, L.,

Espinosa, M., Coll, M., Lorenzo-Diaz, F., and Boer, D.R. (2013). Functional properties and structural requirements of the plasmid pMV158-encoded MobM relaxase domain. *J Bacteriol* 195, 3000-3008 DOI: 10.1128/JB.02264-12.

Fernandez-Lopez, R., Garcillan-Barcia, M.P., Revilla, C., Lazaro, M., Vielva, L., and de la Cruz, F. (2006). Dynamics of the IncW genetic backbone imply general trends in conjugative plasmid evolution. *FEMS Microbiol Rev* 30, 942-966 DOI: 10.1111/j.1574-6976.2006.00042.x.

Figueiredo, T.A., Aguiar, S.I., Melo-Cristino, J., and Ramirez, M. (2006). DNA methylase activity as a marker for the presence of a family of phage-like elements conferring efflux-mediated macrolide resistance in streptococci. *Antimicrob Agents Chemother* 50, 3689-3694 DOI: 10.1128/AAC.00782-06.

Florindo, C., Damiao, V., Silvestre, I., Farinha, C., Rodrigues, F., Nogueira, F., Martins-Pereira, F., Castro, R., Borrego, M.J., Santos-Sanches, I., et al. (2014). Epidemiological surveillance of colonising group B *Streptococcus* epidemiology in the Lisbon and Tagus Valley regions, Portugal (2005 to 2012): emergence of a new epidemic type IV/clonal complex 17 clone. *Euro Surveill* 19

Frost, L.S., Leplae, R., Summers, A.O., and Toussaint, A. (2005). Mobile genetic elements: the agents of open source evolution. *Nat Rev Microbiol* 3, 722-732 DOI: 10.1038/nrmicro1235.

Gawron-Burke, C., and Clewell, D.B. (1982). A transposon in *Enterococcus faecalis* with fertility properties. *Nature* 300, 281-284

Gillings, M.R. (2014). Integrons: past, present, and future. *Microbiol Mol Biol Rev* 78, 257-277 DOI: 10.1128/MMBR.00056-13.

Gyles, C., and Boerlin, P. (2014). Horizontally transferred genetic elements and their role in pathogenesis of bacterial disease. *Vet Pathol* 51, 328-340 DOI: 10.1177/0300985813511131.

Hadjirin, N.F., Harrison, E.M., Holmes, M.A., and Paterson, G.K. (2014). Conjugative transfer frequencies of *mef(A)*-containing Tn1207.3 to macrolide-susceptible *Streptococcus pyogenes* belonging to different *emm* types. *Lett Appl Microbiol* 58, 299-302 DOI: 10.1111/lam.12213.

Henderson-Begg, S.K., Roberts, A.P., and Hall, L.M. (2009). Diversity of putative Tn5253-like elements in *Streptococcus pneumoniae*. *Int J Antimicrob Agents* 33, 364-367 DOI: 10.1016/j.ijantimicag.2008.10.002.

Hernandez-Arriaga, A.M., Espinosa, M., and del Solar, G. (2012). Fitness of the pMV158 replicon in *Streptococcus pneumoniae*. *Plasmid* 67, 162-166 DOI: 10.1016/j.plasmid.2012.01.005.

Iannelli, F., Santagati, M., Santoro, F., Oggioni, M.R., Stefani, S., and Pozzi, G. (2014a). Nucleotide sequence of conjugative prophage Phi1207.3 (formerly Tn1207.3) carrying the *mef(A)/msr(D)* genes for efflux resistance to macrolides in *Streptococcus pyogenes*. *Front Microbiol* 5, 687 DOI: 10.3389/fmicb.2014.00687.

Iannelli, F., Santoro, F., Oggioni, M.R., and Pozzi, G. (2014b). Nucleotide sequence analysis of integrative conjugative element Tn5253 of *Streptococcus pneumoniae*. *Antimicrob Agents Chemother* 58, 1235-1239 DOI: 10.1128/AAC.01764-13.

Jensen, A., and Kilian, M. (2012). Delineation of *Streptococcus dysgalactiae*, its subspecies, and its clinical and phylogenetic relationship to *Streptococcus*

pyogenes. J Clin Microbiol 50, 113-126 DOI: 10.1128/JCM.05900-11.

Johnson, C.M., and Grossman, A.D. (2015). Integrative and conjugative elements (ICEs): what they do and how they work. Annu Rev Genet 49, 577-601 DOI: 10.1146/annurev-genet-112414-055018.

Johnston, C., Martin, B., Fichant, G., Polard, P., and Claverys, J.P. (2014). Bacterial transformation: distribution, shared mechanisms and divergent control. Nat Rev Microbiol 12, 181-196 DOI: 10.1038/nrmicro3199.

Kayansamruaj, P., Pirarat, N., Kondo, H., Hirono, I., and Rodkhum, C. (2015). Genomic comparison between pathogenic *Streptococcus agalactiae* isolated from Nile tilapia in Thailand and fish-derived ST7 strains. Infect Genet Evol 36, 307-314 DOI: 10.1016/j.meegid.2015.10.009.

Kilic, A.O., Vijayakumar, M.N., and al-Khalidi, S.F. (1994). Identification and nucleotide sequence analysis of a transfer-related region in the streptococcal conjugative transposon Tn5252. J Bacteriol 176, 5145-5150

Korona-Glowniak, I., Maj, M., Siwec, R., Niedzielski, A., and Malm, A. (2016). Molecular epidemiology of *Streptococcus pneumoniae* Isolates from children with recurrent upper respiratory tract infections. PLoS One 11, e0158909 DOI:10.1371/journal.pone.0158909.

Korona-Glowniak, I., Siwec, R., and Malm, A. (2015). Resistance determinants and their association with different transposons in the antibiotic-resistant *Streptococcus pneumoniae*. Biomed Res Int 2015, 836496 DOI: 10.1155/2015/836496.

Kowlessar, P.I., O'Connell, N.H., Mitchell, R.D., Elliott, S., and Elliott, T.S. (2006).

Management of patients with *Streptococcus milleri* brain abscesses. J Infect 52, 443-450 DOI: 10.1016/j.jinf.2005.08.028.

Kuroda, M., Ohta, T., Uchiyama, I., Baba, T., Yuzawa, H., Kobayashi, I., Cui, L., Oguchi, A., Aoki, K., Nagai, Y., et al. (2001). Whole genome sequencing of methicillin-resistant *Staphylococcus aureus*. Lancet 357, 1225-1240

Lancefield, R.C. (1933). A serological differentiation of human and other groups of hemolytic Streptococci. J Exp Med 57, 571-595

Lawley, T.D., Klimke, W.A., Gubbins, M.J., and Frost, L.S. (2003). F factor conjugation is a true type IV secretion system. FEMS Microbiol Lett 224, 1-15 DOI: 10.1016/S0378-1097(03)00430-0.

LeBlanc, D.J., Hawley, R.J., Lee, L.N., and St Martin, E.J. (1978). "Conjugal" transfer of plasmid DNA among oral Streptococci. Proc Natl Acad Sci USA 75, 3484-3487

Lekunberri, I., Subirats, J., Borrego, C.M., and Balcazar, J.L. (2017). Exploring the contribution of bacteriophages to antibiotic resistance. Environ Pollut 220, 981-984 DOI: 10.1016/j.envpol.2016.11.059.

Li, Y., Tomita, H., Lv, Y., Liu, J., Xue, F., Zheng, B., and Ike, Y. (2011). Molecular characterization of *erm*(B)- and *mef*(E)-mediated erythromycin-resistant *Streptococcus pneumoniae* in China and complete DNA sequence of Tn2010. J Appl Microbiol 110, 254-265 DOI: 10.1111/j.1365-2672.2010.04875.x.

Li, Y., Yang, L., Fu, J., Yan, M., Chen, D., and Zhang, L. (2017). Microbial pathogenicity and virulence mediated by integrons on Gram-positive microorganisms. Microb Pathog 111, 481-486 DOI:

10.1016/j.micpath.2017.09.035.

Lu, B., Fang, Y., Huang, L., Diao, B., Du, X., Kan, B., Cui, Y., Zhu, F., Li, D., and Wang, D. (2016). Molecular characterization and antibiotic resistance of clinical *Streptococcus dysgalactiae* subsp. *equisimilis* in Beijing, China. *Infect Genet Evol* 40, 119-125 DOI: 10.1016/j.meegid.2016.01.030.

Lybarger, S.R., and Sandkvist, M. (2004). Microbiology. A hitchhiker's guide to type IV secretion. *Science* 304, 1122-1123 DOI: 10.1126/science.1098806.

Mahillon, J., and Chandler, M. (1998). Insertion sequences. *Microbiol Mol Biol Rev* 62, 725-774

Marimon, J.M., Valiente, A., Ercibengoa, M., Garcia-Arenzana, J.M., and Perez-Trallero, E. (2005). Erythromycin resistance and genetic elements carrying macrolide efflux genes in *Streptococcus agalactiae*. *Antimicrob Agents Chemother* 49, 5069-5074 DOI: 10.1128/AAC.49.12.5069-5074.2005.

Mazel, D. (2006). Integrons: agents of bacterial evolution. *Nat Rev Microbiol* 4, 608-620 DOI: 10.1038/nrmicro1462.

McDougal, L.K., Tenover, F.C., Lee, L.N., Rasheed, J.K., Patterson, J.E., Jorgensen, J.H., and LeBlanc, D.J. (1998). Detection of Tn917-like sequences within a Tn916-like conjugative transposon (Tn3872) in erythromycin-resistant isolates of *Streptococcus pneumoniae*. *Antimicrob Agents Chemother* 42, 2312-2318

Menon, T. (2016). Understanding the viridians group Streptococci: Are we there yet? *Indian J Med Microbiol* 34, 421-426 DOI: 10.4103/0255-0857.195371.

Messier, N., and Roy, P.H. (2001). Integron integrases possess a unique

additional domain necessary for activity. *J Bacteriol* 183, 6699-6706 DOI: 10.1128/JB.183.22.6699-6706.2001.

Mingoia, M., Morici, E., Brenciani, A., Giovanetti, E., and Varaldo, P.E. (2014a). Genetic basis of the association of resistance genes *mef(I)* (macrolides) and *catQ* (chloramphenicol) in *Streptococci*. *Front Microbiol* 5, 747 DOI: 10.3389/fmicb.2014.00747.

Mingoia, M., Morici, E., Morroni, G., Giovanetti, E., Del Grosso, M., Pantosti, A., and Varaldo, P.E. (2014b). Tn5253 family integrative and conjugative elements carrying *mef(I)* and *catQ* determinants in *Streptococcus pneumoniae* and *Streptococcus pyogenes*. *Antimicrob Agents Chemother* 58, 5886-5893 DOI: 10.1128/AAC.03638-14.

Mingoia, M., Morici, E., Tili, E., Giovanetti, E., Montanari, M.P., and Varaldo, P.E. (2013). Characterization of Tn5801.Sag, a variant of *Staphylococcus aureus* Tn916 family transposon Tn5801 that is widespread in clinical isolates of *Streptococcus agalactiae*. *Antimicrob Agents Chemother* 57, 4570-4574 DOI: 10.1128/AAC.00521-13.

Mingoia, M., Tili, E., Manso, E., Varaldo, P.E., and Montanari, M.P. (2011). Heterogeneity of Tn5253-like composite elements in clinical *Streptococcus pneumoniae* isolates. *Antimicrob Agents Chemother* 55, 1453-1459 DOI: 10.1128/AAC.01087-10.

Mingoia, M., Vecchi, M., Cochetti, I., Tili, E., Vitali, L.A., Manzin, A., Varaldo, P.E., and Montanari, M.P. (2007). Composite structure of *Streptococcus pneumoniae* containing the erythromycin efflux resistance gene *mef(I)* and the chloramphenicol resistance gene *catQ*. *Antimicrob Agents Chemother* 51, 3983-

3987 DOI: 10.1128/AAC.00790-07.

Noguchi, S., Yatera, K., Kawanami, T., Yamasaki, K., Naito, K., Akata, K., Shimabukuro, I., Ishimoto, H., Yoshii, C., and Mukae, H. (2015). The clinical features of respiratory infections caused by the *Streptococcus anginosus* group. BMC Pulm Med 15, 133 DOI: 10.1186/s12890-015-0128-6.

Ojo, K.K., Ruehlen, N.L., Close, N.S., Luis, H., Bernardo, M., Leitao, J., and Roberts, M.C. (2006). The presence of a conjugative Gram-positive Tn2009 in Gram-negative commensal bacteria. J Antimicrob Chemother 57, 1065-1069 DOI: 10.1093/jac/dkl094.

Palmieri, C., Mingoia, M., Massidda, O., Giovanetti, E., and Varaldo, P.E. (2012). *Streptococcus pneumoniae* transposon Tn1545/Tn6003 changes to Tn6002 due to spontaneous excision in circular form of the *erm(B)*- and *aphA3*-containing macrolide-aminoglycoside-streptothricin (MAS) element. Antimicrob Agents Chemother 56, 5994-5997 DOI: 10.1128/AAC.01487-12.

Parks, T., Barrett, L., and Jones, N. (2015). Invasive streptococcal disease: a review for clinicians. Br Med Bull 115, 77-89 DOI: 10.1093/bmb/ldv027.

Partridge, S.R. (2011). Analysis of antibiotic resistance regions in Gram-negative bacteria. FEMS Microbiol Rev 35, 820-855 DOI: 10.1111/j.1574-6976.2011.00277.x.

Partridge, S.R., Recchia, G.D., Scaramuzzi, C., Collis, C.M., Stokes, H.W., and Hall, R.M. (2000). Definition of the *attI1* site of class 1 integrons. Microbiology 146, 2855-2864 DOI: 10.1099/00221287-146-11-2855.

Perry, J.A., Westman, E.L., and Wright, G.D. (2014). The antibiotic resistome:

what's new? *Curr Opin Microbiol* 21, 45-50 DOI: 10.1016/j.mib.2014.09.002.

Poyart, C., Quesne, G., Acar, P., Berche, P., and Trieu-Cuot, P. (2000). Characterization of the Tn916-like transposon Tn3872 in a strain of *Abiotrophia defectiva* (*Streptococcus defectivus*) causing sequential episodes of endocarditis in a child. *Antimicrob Agents Chemother* 44, 790-793

Pozzi, G., Iannelli, F., Oggioni, M.R., Santagati, M., and Stefani, S. (2004). Genetic elements carrying macrolide efflux genes in Streptococci. *Curr Drug Targets Infect Disord* 4, 203-206

Quintero, B., Araque, M., van der Gaast-de Jongh, C., and Hermans, P.W. (2011). Genetic diversity of Tn916-related transposons among drug-resistant *Streptococcus pneumoniae* isolates colonizing healthy children in Venezuela. *Antimicrob Agents Chemother* 55, 4930-4932 DOI: 10.1128/AAC.00242-11.

Ramos, V., Duarte, C., Diaz, A., and Moreno, J. (2014). Mobile genetic elements associated with erythromycin-resistant isolates of *Streptococcus pneumoniae* in Colombia. *Biomedica* 34 Suppl 1, 209-216 DOI: 10.1590/S0120-41572014000500023.

Rato, M.G., Bexiga, R., Nunes, S.F., Vilela, C.L., and Santos-Sanches, I. (2010). Human group A Streptococci virulence genes in bovine group C Streptococci. *Emerg Infect Dis* 16, 116-119 DOI: 10.3201/eid1601.090632.

Remington, L.T., and Sligl, W.I. (2014). Community-acquired pneumonia. *Curr Opin Pulm Med* 20, 215-224 DOI: 10.1097/MCP.0000000000000052.

Roberts, A.P., and Mullany, P. (2009). A modular master on the move: the Tn916 family of mobile genetic elements. *Trends Microbiol* 17, 251-258 DOI:

10.1016/j.tim.2009.03.002.

Roberts, A.P., and Mullany, P. (2011). Tn916-like genetic elements: a diverse group of modular mobile elements conferring antibiotic resistance. *FEMS Microbiol Rev* 35, 856-871 DOI: 10.1111/j.1574-6976.2011.00283.x.

Rolo, D., Fenoll, A., Ardanuy, C., Calatayud, L., Cubero, M., de la Campa, A.G., and Linares, J. (2011). Trends of invasive serotype 6C pneumococci in Spain: emergence of a new lineage. *J Antimicrob Chemother* 66, 1712-1718 DOI: 10.1093/jac/dkr193.

Sadowy, E., Matynia, B., and Hryniewicz, W. (2010). Population structure, virulence factors and resistance determinants of invasive, non-invasive and colonizing *Streptococcus agalactiae* in Poland. *J Antimicrob Chemother* 65, 1907-1914 DOI: 10.1093/jac/dkq230.

Santagati, M., Iannelli, F., Cascone, C., Campanile, F., Oggioni, M.R., Stefani, S., and Pozzi, G. (2003). The novel conjugative transposon Tn1207.3 carries the macrolide efflux gene *mef(A)* in *Streptococcus pyogenes*. *Microb Drug Resist* 9, 243-247 DOI: 10.1089/107662903322286445.

Santagati, M., Iannelli, F., Oggioni, M.R., Stefani, S., and Pozzi, G. (2000). Characterization of a genetic element carrying the macrolide efflux gene *mef(A)* in *Streptococcus pneumoniae*. *Antimicrob Agents Chemother* 44, 2585-2587

Santagati, M., Lupo, A., Scillato, M., Di Martino, A., and Stefani, S. (2009). Conjugal mobilization of the mega element carrying *mef(E)* from *Streptococcus salivarius* to *Streptococcus pneumoniae*. *FEMS Microbiol Lett* 290, 79-84 DOI: 10.1111/j.1574-6968.2008.01408.x.

Santoro, F., Oggioni, M.R., Pozzi, G., and Iannelli, F. (2010). Nucleotide sequence and functional analysis of the *tet(M)*-carrying conjugative transposon Tn5251 of *Streptococcus pneumoniae*. FEMS Microbiol Lett 308, 150-158 DOI: 10.1111/j.1574-6968.2010.02002.x.

Schroder, G., Krause, S., Zechner, E.L., Traxler, B., Yeo, H.J., Lurz, R., Waksman, G., and Lanka, E. (2002). TraG-like proteins of DNA transfer systems and of the *Helicobacter pylori* type IV secretion system: inner membrane gate for exported substrates? J Bacteriol 184, 2767-2779

Scott, K.P. (2002). The role of conjugative transposons in spreading antibiotic resistance between bacteria that inhabit the gastrointestinal tract. Cell Mol Life Sci 59, 2071-2082

Sengupta, M., and Austin, S. (2011). Prevalence and significance of plasmid maintenance functions in the virulence plasmids of pathogenic bacteria. Infect Immun 79, 2502-2509 DOI: 10.1128/IAI.00127-11.

Shi, L., Zheng, M., Xiao, Z., Asakura, M., Su, J., Li, L., and Yamasaki, S. (2006). Unnoticed spread of class 1 integrons in Gram-positive clinical strains isolated in Guangzhou, China. Microbiol Immunol 50, 463-467

Shoemaker, N.B., Smith, M.D., and Guild, W.R. (1979). Organization and transfer of heterologous chloramphenicol and tetracycline resistance genes in pneumococcus. J Bacteriol 139, 432-441

Siguié, P., Gourbeyre, E., and Chandler, M. (2014). Bacterial insertion sequences: their genomic impact and diversity. FEMS Microbiol Rev 38, 865-891 DOI: 10.1111/1574-6976.12067.

Smillie, C., Garcillan-Barcia, M.P., Francia, M.V., Rocha, E.P., and de la Cruz, F. (2010). Mobility of plasmids. *Microbiol Mol Biol Rev* 74, 434-452 DOI: 10.1128/MMBR.00020-10.

Stewart, P.E., Byram, R., Grimm, D., Tilly, K., and Rosa, P.A. (2005). The plasmids of *Borrelia burgdorferi*: essential genetic elements of a pathogen. *Plasmid* 53, 1-13 DOI: 10.1016/j.plasmid.2004.10.006.

Storrs, M.J., Poyart-Salmeron, C., Trieu-Cuot, P., and Courvalin, P. (1991). Conjugative transposition of Tn916 requires the excisive and integrative activities of the transposon-encoded integrase. *J Bacteriol* 173, 4347-4352

Sui, Z., Zhou, W., Yao, K., Liu, L., Zhang, G., Yang, Y., and Feng, J. (2013). Complete genome sequence of *Streptococcus pneumoniae* strain A026, a clinical multidrug-resistant isolate carrying Tn2010. *Genome Announc* 1 DOI: 10.1128/genomeA.01034-13.

Swartley, J.S., McAllister, C.F., Hajjeh, R.A., Heinrich, D.W., and Stephens, D.S. (1993). Deletions of Tn916-like transposons are implicated in *tetM*-mediated resistance in pathogenic *Neisseria*. *Mol Microbiol* 10, 299-310 DOI: 10.1111/j.1365-2958.1993.tb01956.x.

Takamatsu, D., Osaki, M., and Sekizaki, T. (2003). Chloramphenicol resistance transposable element TnSs1 of *Streptococcus suis*, a transposon flanked by IS6-family elements. *Plasmid* 49, 143-151

Talebi, M., Azadegan, A., Sadeghi, J., Ahmadi, A., Ghanei, M., Katouli, M., Owlia, P., and Pourshafie, M.R. (2016). Determination of characteristics of erythromycin resistant *Streptococcus pneumoniae* with preferred PCV usage in Iran. *PLoS One* 11, e0167803 DOI: 10.1371/journal.pone.0167803.

- Tantivitayakul, P., Lapirattanakul, J., Vichayanrat, T., and Muadchiengka, T. (2016). Antibiotic resistance patterns and related mobile genetic elements of Pneumococci and beta-Hemolytic Streptococci in Thai Healthy Children. *Indian J Microbiol* 56, 417-425 DOI: 10.1007/s12088-016-0607-3.
- Tomich, P.K., An, F.Y., and Clewell, D.B. (1980). Properties of erythromycin-inducible transposon Tn917 in *Streptococcus faecalis*. *J Bacteriol* 141, 1366-1374
- Varaldo, P.E., Montanari, M.P., and Giovanetti, E. (2009). Genetic elements responsible for erythromycin resistance in Streptococci. *Antimicrob Agents Chemother* 53, 343-353 DOI: 10.1128/AAC.00781-08.
- Verani, J.R., McGee, L., Schrag, S.J. (2010). Prevention of perinatal group B Streptococcal disease-revised guidelines from CDC, 2010. *MMWR Recomm Rep* 59, 1-36
- Vijayakumar, M.N., Priebe, S.D., and Guild, W.R. (1986). Structure of a conjugative element in *Streptococcus pneumoniae*. *J Bacteriol* 166, 978-984
- Walker, M.J., Barnett, T.C., McArthur, J.D., Cole, J.N., Gillen, C.M., Henningham, A., Sriprakash, K.S., Sanderson-Smith, M.L., and Nizet, V. (2014). Disease manifestations and pathogenic mechanisms of group A Streptococcus. *Clin Microbiol Rev* 27, 264-301 DOI: 10.1128/CMR.00101-13.
- Wang, H., Roberts, A.P., and Mullany, P. (2000). DNA sequence of the insertional hot spot of Tn916 in the *Clostridium difficile* genome and discovery of a Tn916-like element in an environmental isolate integrated in the same hot spot. *FEMS Microbiol Lett* 192, 15-20 DOI: 10.1111/j.1574-6968.2000.tb09352.x.
- Wright, L.D., and Grossman, A.D. (2016). Autonomous replication of the

conjugative transposon Tn916. J Bacteriol 198, 3355-3366 DOI: 10.1128/JB.00639-16.

Wyres, K.L., van Tonder, A., Lambertsen, L.M., Hakenbeck, R., Parkhill, J., Bentley, S.D., and Brueggemann, A.B. (2013). Evidence of antimicrobial resistance-conferring genetic elements among pneumococci isolated prior to 1974. BMC Genomics 14, 500 DOI: 10.1186/1471-2164-14-500.

Zhang, C., Zhang, Z., Song, L., Fan, X., Wen, F., Xu, S., and Ning, Y. (2015). Antimicrobial resistance profile and genotypic characteristics of *Streptococcus suis* capsular type 2 isolated from clinical carrier sows and diseased pigs in China. Biomed Res Int 2015, 284303 DOI: 10.1155/2015/284303.

Zhou, W., Yao, K., Zhang, G., Yang, Y., Li, Y., Lv, Y., and Feng, J. (2014). Mechanism for transfer of transposon Tn2010 carrying macrolide resistance genes in *Streptococcus pneumoniae* and its effects on genome evolution. J Antimicrob Chemother 69, 1470-1473 DOI: 10.1093/jac/dku019.

